

DEPARTMENT OF MINES AND ENERGY  
SOUTH AUSTRALIA

Rept.Bk.No. 79/19

PRECAMBRIAN AND PALAEOZOIC GEOLOGY  
OF THE PEAKE AND DENISON RANGES

GEOLOGICAL SURVEY

by


G.J. AMBROSE

and

R.B. FLINT

REGIONAL GEOLOGY DIVISION

G.S. No. 6138  
D.M. No. 302/72



<u>CONTENTS</u>	<u>PAGE</u>
ABSTRACT	1
INTRODUCTION	2
GEOGRAPHY	2
PREVIOUS INVESTIGATIONS	5
GEOLOGICAL SETTING OF THE PRECAMBRIAN ROCKS	7
STRATIGRAPHY	9
PEAKE METAMORPHICS (? LOWER PROTEROZOIC)	10
Unnamed metamorphics E d1	11
Tidnamurkuna Volcanics E dt	12
Unnamed schists E d2	13
Baltucoodna Quartzite E db	13
Unnamed metamorphics E d3	15
Correlation	15
WIRRIECURRIE GRANITE E W (CARPENTARIAN)	16
CALLANNA BEDS (WILLOURAN)	18
Younghusband Conglomerate E co	20
Coominaree Dolomite E ck	23
Cadlareena Volcanics E cc	24
SEQUENCES OF UNCERTAIN AGE (WILLOURAN OR TORRENSIAN)	33
Rockwater Beds E ad	35
War Loan Beds E aw	36
Nilpinna Beds E an	36
Duff Creek Beds E af	37
Murrana Beds E aa	40
Undifferentiated Blocks E a	42
BURRA GROUP (TORRENSIAN)	44
Unnamed siltstone E bl	45
Fountain Spring Beds E bl	46
Mount Margaret Quartzite E bo	47
Skillogalee Dolomite E bk	48
Unnamed transitional unit E b2	52
Kalachalpa Formation E bh	55
UMBERATANA GROUP (STURTIAN - MARINOAN)	57
Calthorinna Tillite E ub	58
Unnamed sandstone E ul	63
Tapley Hill Formation E ft	64
Thora Dolomite E fl	65
Unnamed siltstone E hl	65
BUNGADILLINA MONZONITE Odb (ORDOVICIAN)	66
BOORTHANNA FORMATION Plb (PERMIAN)	69
DIAPIRISM	70
TECTONISM	74
METAMORPHISM	77
ECONOMIC GEOLOGY	79
HISTORICAL MINING ACTIVITIES	79
GEOCHEMISTRY	82
SUMMARY	83
ACKNOWLEDGEMENTS	86
REFERENCES	87
APPENDICES	
<u>FIGURES</u>	<u>PHOTO NO or DRAWING NO</u>
Fig. 1. Map of the Peake and Denison Ranges showing local geography.	S13007
Fig. 2. Landsat Imagery, Peake and Denison Ranges and environs.	S13265
Fig. 3. Regional geological setting.	S13259
Fig. 4. Photomicrograph of amygdaloidal basalt, Tidnamurkuna Volcanics.	30296

Fig. 5.	Photomicrograph of plagioclase phenocryst, Tidnamurkuna Volcanics.	30297
Fig. 6.	Hand specimen of porphyritic rhyolite, Tidnamurkuna Volcanics.	30298
Fig. 7.	Plagioclase phenocrysts in basalt, Baltucoodna Quartzite.	30299
Fig. 8.	Heavy-mineral banding in quartzites near Spring Hill.	30300
Fig. 9.	Feldspar phenocrysts, Wirriecurrie Granite.	30301
Fig. 10.	Younghusband Conglomerate and basement/cover unconformity.	30302
Fig. 11.	Cadlareena Volcanics and basement/cover unconformity.	30303
Fig. 12.	Vesicular basalts and lenticular mudstone, Cadlareena Volcanics.	30304
Fig. 13.	AFM diagram, Cadlareena and Wooltana Volcanics.	S13300
Fig. 14.	Cross-laminations in sandstones of the Nilpinna Beds.	30305
Fig. 15.	Laminated dolomites of the Duff Creek Beds.	30306
Fig. 16.	Mud cracks in silty dolomite, Duff Creek Beds.	30307
Fig. 17.	Aerial view of the eastern margin of Margaret Inlier.	30308
Fig. 18.	Aerial view of Mount Margaret Quartzite and Mount Margaret Plateau.	30309
Fig. 19.	Aerial view of Skillogalee Dolomite.	30310
Fig. 20.	Boorthanna geological section.	78-314
Fig. 21.	Intraformational chert pebble conglomerate, unnamed transition unit.	30311
Fig. 22.	Typical intraformational dolomite, upper Burra Group.	30312
Fig. 23.	Columnar stromatolites in the Kalachalpa Formation.	30313
Fig. 24.	Conglomeratic and massive dolomites, Calthorinna Tillite.	30314
Fig. 25.	Conglomeratic dolomite and diamictite, Calthorinna Tillite.	30315
Fig. 26.	Dolomite with reworked diamictite clasts, Calthorinna Tillite.	30316
Fig. 27.	Porphyritic monzonite, Bungadillina Monzonite.	30317
Fig. 28.	Aligned feldspar phenocrysts, Bungadillina Monzonite.	30318
Fig. 29.	Zoned plagioclase phenocryst, Bungadillina Monzonite.	30319
Fig. 30.	Angular unconformity between Permian Boorthanna Formation and Torrensian Kalachalpa Formation.	30320
Fig. 31.	Aerial view of disoriented blocks in diapir.	30321
Fig. 32.	Diapiric, carbonate matrix breccia.	30322

## TABLES

	<u>PAGE</u>
1. Stratigraphic summary, Peake and Denison Ranges.	10A
2. Silicate analyses, basalts and rhyolites, Tidnamurkuna Volcanics and Baltucoodna Quartzite.	14A
3. Silicate analyses, Wirriecurrie Granite.	17
4. Stratigraphy, Callanna Beds.	19
5. Correlation chart for Adelaidean stratigraphy.	20A-76-97
6. Silicate analyses, Cadlareena Volcanics.	30

7.	Trace element analyses, Cadlareena Volcanics.	33
8.	Stratigraphy, sequences of uncertain age.	34
9.	Stratigraphy, Burra Group.	44
10.	Stratigraphy, Umberatana Group.	57
11.	Silicate analyses, Bungadillina Monzonite.	67A
12.	Trace element means for various rock groups.	82A-S13334

## ENCLOSURES

## DRAWING NO

Encl. 1	Geological map, Peake and Denison Ranges.	78-867 & 78-868
Encl. 2	Geological block diagrams, Peake and Denison Ranges.	78-317
Encl. 3	Special Mining Lease and Exploration Licence map.	78-316
Encl. 4	Geochemical values, Peake and Denison Ranges.	78-315

## APPENDICES

A	Definition of New Units	
B	Stratigraphic type sections for new units in the Callanna Beds.	78-211
C	Stratigraphic type sections for new units in sequences of uncertain age.	78-214
D	Stratigraphic type sections for new units in the Burra Group.	78-213
E	Stratigraphic type sections for new units in the Umberatana Group.	78-212



DEPARTMENT OF MINES AND ENERGY  
SOUTH AUSTRALIA

Rept.Bk.No. 79/19  
G.S. No. 6138  
D.M. No. 302/72

PRECAMBRIAN AND PALAEOZOIC GEOLOGY OF  
THE PEAKE AND DENISON RANGES

ABSTRACT

The Peake and Denison Ranges comprise four northerly-trending Precambrian inliers in the southwestern portion of the Great Artesian Basin. The Peake Metamorphics (?Lower Proterozoic) the oldest rocks exposed, form inliers of interlayered metabasalts and quartzites, in sequences over 10 000 m thick. These rocks were subjected to greenschist to amphibolite facies metamorphism and syntectonic granite intrusion about 1600 Ma, and further deformation and probable greenschist facies metamorphism at 1050 Ma.

Unconformably overlying the Peake Metamorphics are at least 29 000 m of Adelaidean paralic sediments deposited in a steadily subsiding, faultbounded trough which was a northern extension of the Adelaide Geosyncline. The thickness (27 000 m) of Willouran and Torrensian sediments preserved here is greater than elsewhere in the geosyncline. Adelaidean sedimentation is first recorded by the Willouran Callanna Beds. A basal conglomerate-quartzite (Younghusband Conglomerate) is overlain by the stromatolitic Coominaree Dolomite and the predominantly basaltic Cadlareena Volcanics. Faulting and diapirism have caused extensive disruption of the sequence above the Cadlareena Volcanics. From the dismembered succession the following stratigraphic units of Willouran or Torrensian age have been recognised; Rockwater Beds, War Loan Beds, Nilpinna Beds, Duff Creek Beds, and Murrana Beds.

The overlying Torrensian sequence (Burra Group) exceeds 10 000 m and comprises; unnamed siltstones, Fountain Spring Beds, Mount Margaret Quartzite, Skillogalee Dolomite, unnamed transition unit, and Kalachalpa Formation.

A period of mild erosion preceeded deposition of the Sturtian Umberatana Group. The onset of glacial conditions resulted in deposition of the glaciomarine Calthorinna Tillite and interfingering sandstones. Subsequent marine silty shales and silty dolomites of the Tapley Hill Formation are overlain by the regressive Thora Dolomite and Willochra Subgroup.

Folding, diapirism and metamorphism (to low greenschist facies) of the Adelaidean succession during the Cambro-Ordovician Delamerian Orogeny was accompanied by late-tectonic intrusion of porphyritic monzonites, syenites, diorites and albitites (Bungadillina Monzonite).

## INTRODUCTION

The Peake and Denison Ranges are approximately 1000 km north of Adelaide. They trend northerly for 130 km parallel to the Central Australia Railway (narrow gauge) and Marree-Oodnadatta road, and cover an area of 2 600 km<sup>2</sup> between latitudes 27°45' and 28°55'S, and longitudes 135°30' and 136°25'E (Figs 1-3).

The first comprehensive map of the area was compiled by Reyner (1955). In 1972 R.P. Coats and W.V. Preiss mapped a portion of the ranges as part of the programme to map WARRINA on which the ranges are mainly situated. Subsequent detailed mapping by the authors during 1975-76 completed this mapping programme. Field observations were plotted on 1:20 000 scale aerial photographs (Department of Lands, Surveys 88, 89, 90). Geological data were transferred onto 1:77 000 aerial photographs (Department of Lands, Surveys 588, 591) and subsequently onto photoscale base maps. From these bases a 1:100 000 geological map was prepared (Encl. 1).

## GEOGRAPHY

The Peake and Denison Ranges consist of a northerly trending series of low-lying ranges with a maximum relief of 300 m above vast undulating plains. Four Precambrian inliers constitute the Peake and Denison Ranges - the Dutton, Algebuckina, Denison and Margaret Inliers. The Dutton and Algebuckina Inliers lie to the north of and are subordinate to the Denison and Margaret Inliers which form the bulk of the ranges (Figs 1 & 2).

The southernmost range (Margaret Inlier) extends northward from Anna Creek H.S. for 70 km and averages 16 km in width. This inlier is named after Mount Margaret (410 m a.s.l.) the highest peak in the region. Topography is particularly rugged along the eastern margin of the inlier. The Mount Margaret Plateau, a conspicuous, gently dipping peneplain, comprises the northern core of the Margaret Inlier. This imposing plateau rises about 250 m above the surrounding plains and from its maximum elevation just north of Mount Margaret it slopes gently westward and northward.

The Denison Inlier rises abruptly from the plains about 10 km north of the Margaret Inlier and extends northward for about 25 km with an average width of 7 km. Mounts Denison and Kingston are two high points in the inlier. Approximately 7 km northwest of the Denison Inlier, on the banks of the Neales River, are a series of small low hills comprising the Algebuckina Inlier. The Dutton Inlier is named after Mount Dutton, the highest point in a small prominent range of hills about 15 km northwest of Algebuckina Inlier.

Rock exposures are good except on the western margin of the ranges where Mesozoic weathering, leaching and kaolinization of Precambrian rocks has resulted in a subdued topography. Mesozoic and Permian sediments lap onto the western margin of the ranges and weathered Precambrian rocks are often masked by a veneer of younger sediments and associated lag deposits.

Drainage from the ranges is dominantly easterly with creeks draining via the Peake Creek and Neales River into Lake Eyre 80 km to the east. There are two permanent waterholes in the

region, one near Algebuckina R.S. on the Neales River and the other at Warrawaroon W.H. on the Peake Creek. Along the faulted eastern margin of the ranges are several springs which include Freeling, Edith and Tarlton Springs. On the plains to the east of the ranges are numerous artesian springs (e.g. Hawker, Outside and Fanny Springs) and artesian bores (e.g. Hope Creek, Birthday and Umbum Bores - see Encl. 1).

Climate of the region is arid with a mean annual rainfall of 125 mm and average annual evaporation of 3800 mm. The low rainfall falls erratically mainly from thunderstorms between October and May. The region experiences hot summers with an average summer maximum temperature of  $36.5^{\circ}\text{C}$  while minimum winter temperatures often fall below  $0^{\circ}\text{C}$ .

Although vegetation in the region is sparse a considerable variety of species is present (Laut et al., 1977). The ranges themselves are hosts to Acacia aneura (mulga), A. brachystachya (umbrella mulga), Eremophila freelingii, Digitaria browaii (cotton grass), Bassia spp. (bindyi) and Eremophila maculata (native fuchsia). Areas marginal to the ranges contain varying proportions of the following species; Atriplex rhagodioides (silver saltbush), A. vesicaria (bladder saltbush), Maireana astrotricha, Cucumis myriocarpies (paddy melon), Kennedia astrotricha (pearl bush), Salicornia Spp. (samphire) and bindyi. The following species are found in dune country on the plains; Acacia ligulata (sandhill wattle), Zygochloa paradoxa (sandhill cane grass), Cassia spp. (Kangaroo bush), Eremophila spp., Dodonaea attenuata (hop bush), native fuchsia and mulga. Large trees are restricted to creeks draining from the ranges and species include Eucalyptus microtheca (coolibah), E. camaldulensis (river red gum) and Acacia cambagei (gidgee).

Because of the harsh climate the region is sparsely settled, and since the area lies north of (outside) the dingo-proof "Dog-

Fence" land use is restricted to cattle grazing. The four cattle stations are Anna Creek, Peake, Nilpinna and Allendale. A very small township is located at William Creek and Commonwealth Railway settlements at Mount Dutton, Edwards Creek and Anna Creek.

The Central Australia Railway (narrow gauge) and graded Marree-Oodnadatta road provide reasonable access to the region. Through the ranges an erratic system of rough tracks allow access, the adequacy of which varies according to weather conditions.

#### PREVIOUS INVESTIGATIONS

John McDouall Stuart, who explored much of Central Australia in the late 1850's and early 1860's, made the first recorded geological observations in the Peake and Denison Ranges (Stuart, 1860; 1862). Scouler (1885) described the topography from Anna Creek to Mount Margaret, analysed water samples from several bores and wells, and described quartzites and sandstones in the area. Basement metamorphics near Mount Denison, granites and gneisses near Mount Kingston, and a quartz syenite at Spring Hill were described by East (1889). Shortly after, Brown (1894) found traces of gold and a 1-carat diamond while panning on the western side of the ranges near Peake Creek.

Copper prospecting was active in the late 1890's after the establishment of a permanent settlement at "Peake", a station on the original overland telegraph line from Adelaide to Darwin. A smelting plant was built near the telegraph station to treat ore from the Peake and Coppertop Mines. In the period 1915-1920 copper was mined from the War Loan, Coominaree and Last Chance Mines and early mining activities at these localities were reported by Winton (1919).

Preliminary observations on the geology and hydrology of the area were made by Jack (1930). The first detailed hydrological

report was written by Chugg (1957). Regional geological mapping of the Peake and Denison Ranges during 1953 resulted in the publication of seven 1:63 360 geological maps, namely Algebuckina, Nilpinna, Conway, Umbum, Boorthanna, Cadlareena and Anna (Dickinson et al., 1954 a; 1954b; 1955a; 1955b; 1955c; 1955d; 1955e). The first comprehensive study of the geology of the region was published by Reyner in 1955. He divided the Adelaidean sequence into the following units:

Top Sturtian - The Sturtian Tillite (1300m)

Torrensian - the "Magnesite Series" (3600m)

- the Mount Margaret Quartzites (3200m)

Willouran - the Duff Creek Formation (3200m)

In 1964, Thomson and Coats proposed the name Duff Creek Beds to replace Duff Creek Formation which they correlated with the type section of the Callanna Beds in the Willouran Ranges, 240 km southeast of the Peake and Denison Ranges. The Mount Margaret Formation (a modification of Reyner's term) was considered equivalent to the Witchelina and Copley Quartzites, then believed to be at the base of the Burra Group.

In 1966, Thomson described a basal Adelaidean (Willouran) sequence unconformably overlying crystalline basement and in faulted contact with younger Duff Creek Beds. From these oldest Adelaidean rocks, Preiss (1973) has reported columnar stromatolites. Schopf and Fairchild (1973) and Fairchild (1975) have studied late Precambrian stromatolites and microbiota and their stratigraphic significance.

Detailed stratigraphic sections through Adelaidean sequences within the ranges and Mesozoic rocks flanking the ranges were first measured in 1963 by French Petroleum Company geologists (Cooper et al., 1963).

Since 1920 no mines have been worked; however, in the past decade mineral exploration has been carried out for U, Cu, Pb, Zn, Ni, Co, Cr, Mn, Ag, As, and Au in the Precambrian rocks and U, Au, S, K, clay, coal and liquid hydrocarbons in the younger sedimentary basins flanking the ranges (Encl. 3). In the period 1966-1968 North Broken Hill Ltd carried out extensive soil and stream sediment sampling and drilled three diamond drillholes, Peake 1 and War Loan 1 and 2 (Forwood, 1968). In 1970 Australasian Mining Corporation took out a special mining lease and conducted radiometric and aeromagnetic surveys, and drilled three percussion and four diamond drillholes, Algebuckina 1, 2, 3 and 4 (Sargeant, 1970). Geochemical and geophysical investigations of copper-uranium mineralisation at the Last Chance Mine indicated no economic mineralisation (Leeson, 1972). Uranerz (Australia) Pty Ltd explored for uranium and drilled two diamond drill holes, one in the Cadlareena Volcanics (2BAA/DDHI) and the other in migmatites of the Peake Metamorphics (DDH2/MG2) - (Iliff et al., 1974; Iliff, 1975). Recently, Western Mining Corporation Ltd carried out extensive stream sediment surveys and soil and rock chip sampling for base metals (Western Mining Corporation Ltd, 1975).

#### GEOLOGICAL SETTING OF THE PRECAMBRIAN ROCKS

The Peake and Denison Ranges are a series of northerly-trending Precambrian inliers situated in the southwestern portion of the Great Artesian Basin, approximately midway between the Musgrave Block and Willouran Ranges (Fig. 3). They are linked to the Willouran Ranges and Mount Painter Province by the Muloorina Gravity High, and consist of ? Lower Proterozoic Peake Metamorphics unconformably overlain by Adelaidean sediments. The Peake Metamorphics have no apparent correlatives in South Australia and are tentatively correlated with the Eastern Creek Volcanics of the Mount Isa Trough. Adelaidean sedimentation represents

deposition in the northern extension of the Adelaide Geosyncline which was intermittently connected to other basins to the north and east (i.e. Amadeus Basin and Broken Hill area).

The Peake and Denison Ranges are flanked by two Permian basins - the Arckaringa Basin to the west and Pedirka Basin to the northeast. Beneath the Boorthanna Trough (a sub-basin of the Arckaringa Basin) early Palaeozoic sediments intersected in drill holes include ? Ordovician quartzites in Weedina No. 1 (Papalia, 1970) and Devonian dolomites, arenites, and anhydrite of the Cootanoorina Formation in Cootanoorina No. 1 and Weedina No. 1 (Papalia, 1970; Allchurch et al., 1973). Below the Pedirka Basin a more complete early Palaeozoic sequence is represented comprising Cambrian carbonates and shales, Ordovician sands and Devonian terrestrial sands, shales and conglomerates (Youngs, 1975a).

The Permian sediments of the Arckaringa and Pedirka Basins have received considerable attention in recent years (Ludbrook, 1961; Holmes, 1970; Wopfner, 1970; Holmes and Rayment, 1970; Papalia, 1970; Allchurch et al., 1973; Townsend, 1973; Barclay, 1974; Townsend and Ludbrook, 1975; Youngs, 1975a; 1975b). In the Boorthanna Trough, the base of the Permian is marked by glacial diamictites, conglomerates, pebbly sandstones and grey clays of the Boorthanna Formation. Diamictites and conglomerates of this basal unit outcrop along the western margin of the ranges. The Boorthanna Formation is overlain by marine shales of the Stuart Range Formation which is in turn overlain by the Mount Toondina Formation, a terrestrial sequence of coals, carbonaceous shales and siltstones, and arenaceous sediments (Townsend and Ludbrook, 1975).



In the Pedirka Basin, glacial diamictites, pebbly sandstones and clays of the Crown Point Formation are overlain by the Purni Formation, a terrestrial sequence of sandstones, siltstones, shales, conglomerates and thin coal bands (Youngs, 1975 a; 1975b).

Great Artesian Basin sediments flanking the ranges consist of ? Jurassic basal conglomeratic, kaolinitic sandstones (Algebuckina Sandstone) and overlying Cretaceous micaceous and arenaceous sediments of the Cadna-owie Formation (Wopfner & Heath, 1963; Wopfner et al., 1970). Other Cretaceous sediments include carbonaceous shales and concretionary, fossiliferous limestones of the Bulldog Shale (Freytag, 1966; Teluk, 1974) and argillaceous siltstones and shales, and olive glauconitic sandstones of the Oodnadatta Formation (Freytag et al., 1967).

Tertiary sediments comprise silicified claystones and sandstones, and silcrete conglomerates (Freytag et al., 1967). They have been extensively eroded with pebbles, cobbles and boulders of silicified sediments occurring as a Quaternary gibber lag over Mesozoic units. Rare outcrops of Tertiary sediments have been preserved at Mount Harvey and along the eastern margin of the ranges.

During the late Miocene and Pleistocene faulting resulted in uplift of huge blocks tilted gently to the west, forming the Peake and Denison Ranges (Wopfner, 1968; 1972).

Quaternary sediments surrounding the ranges include sand dunes, gibber plains, calcareous silts and clays, gypsiferous gravels and alluvial sands, silts and clays.

#### STRATIGRAPHY

Geological mapping by Reyner (1955) and Dickinson et al. (1954a; 1954b; 1955a; 1955b; 1955c; 1955d; 1955e) established the outcrop boundaries and general stratigraphy for the ranges.

However the stratigraphic succession and facing for the Peake Metamorphics were not known and the existence and extent of volcanics were not fully realised. The Adelaidean succession was only partly assembled. The structures now recognised as diapirs were interpreted as complex crush zones resulting from interaction of thrusting, rifting and polyphase folding. During the present mapping programme these zones were remapped in detail and are shown on the geological map (Encl. 1). Details of the revised stratigraphic succession are highlighted in Table 1.

#### PEAKE METAMORPHICS (?LOWER PROTEROZOIC)

Pre-Adelaidean basement rocks chiefly outcrop in the northern portion of the Peake and Denison Ranges (Encl. 1). Reyner (1955) originally named the basement rocks the Peake Series which was subsequently amended to Peake Metamorphics (Thomson, 1966). However an intrusive granite (Wirriecurrie Granite) has been recognised and it is proposed that the term Peake Metamorphics exclude the Wirriecurrie Granite and be restricted to the remainder of the pre-Adelaidean rocks.

Metamorphics can be subdivided into five mappable units (Table 1).

- a. Unnamed metamorphics (Pd1). This unit consists of a wide variety of rock types including quartzites, quartz-biotite and muscovite schists, quartz-biotite-feldspar gneisses, migmatites, pegmatites and diorites.
- b. Tidnamurkuna Volcanics (Bdt). Volcanics within this unit include flow-banded porphyritic rhyolites and amygdaloidal basalts with a 600 m section exposed.
- c. Unnamed schists (Pd2). Rock types are dominantly quartz, muscovite schists and quartz, chlorite phyllites with minor sandstones.

- d. Baltucoodna Quartzite (Edb). The unit is characterised by interlayered greyish-white quartzites and basalts; and minor sillimanite gneisses, pyroxene granulites, calcite marbles and hornblende, epidote calc-silicates.
- e. Unnamed metamorphics (Ed3). A wide variety of rock types present include grey quartzites, magnetite-rich epidiosites and arenaceous schists, minor clinopyroxene granulites and amphibolites.

The Baltucoodna Quartzite conformably overlies the unnamed schists, which in turn conformably overlie the Tidnamurkuna Volcanics. However, the stratigraphic relationships of unnamed metamorphics (Ed1 and Ed3) with themselves and the other units are not known.

(a) Unnamed metamorphics Ed1

Metamorphic rocks outcrop in the Algebuckina Inlier and in the northern end of the Denison Inlier. In the Algebuckina Inlier garnetiferous quartz-biotite schists and quartz-feldspar-biotite gneisses are common. The garnets are red, euhedral and up to 10 mm in size. Quartz-biotite metasandstones and quartzites are rare while diorite and pegmatites (with coarse, graphic quartz-feldspar intergrowths) are common.

In the Denison Inlier, outcropping metamorphics are quartz-biotite and quartz-muscovite schists, quartz-feldspar-biotite and plagioclase-hornblende gneisses and greyish-white quartzites. Ellipsoidal pods of epidote occur sporadically within the quartzites. Dioritic and pegmatitic sills and dykes are common. Metamorphosed basalts could be widespread as rocks containing amygdalae with quartz centres and radiating, actinolite-rich margins were observed in several localities: however, metamorphism and a strong cleavage renders distinction between diorites and basic volcanics difficult.

Migmatites are locally developed just north of Peake Creek and indicate middle to upper amphibolite facies metamorphism.

K-Ar dating on hornblende from two diorite sills yielded ages of  $1508 \pm 30$  Ma and  $1453 \pm 30$  Ma (Webb, 1974), recording the metamorphic age for the area.

(b) Tidnamurkuna Volcanics Edt

The Tidnamurkuna Volcanics consist of greater than 600 m of amygdaloidal basalts, porphyritic rhyolites, epidiosites and minor phyllites which outcrop southwest of Peake ruins and north of Mount Denison. The sequence is dominantly basaltic with two major periods of acidic eruptions.

The common minerals within the basalts are actinolite, hornblende, plagioclase, epidote, chlorite and magnetite. Plagioclase laths (up to 0.5 mm and very rarely to 10 mm in length) show only poorly developed polysynthetic twinning and exhibit a decussate texture. Amygdales are common and have been filled by quartz and radiating aggregates of fibrous epidote crystals (Fig. 4).

A lower, single rhyolite flow up to 5 m thick has plagioclase, potash feldspar and quartz phenocrysts. Phenocryst abundance varies from 10 to 30% with subhedral to ovoid feldspar (up to 20 mm) and spherical bluish-grey quartz phenocrysts averaging 2 mm in size (Figs. 5 & 6). Quartz phenocrysts are common near (but not restricted to) the margins while very coarse feldspar phenocrysts occur in the centre of the flow.

An upper rhyolitic unit contains about 250 m of interlayered rhyolites, epidiosites and basalts. These rhyolites are also strongly porphyritic with phenocrysts of potash feldspar,

plagioclase and quartz. Phenocrysts often show undulose extinction or have been completely recrystallised forming elongate aggregates of smaller grains. The aphanitic ( 0.4 mm) groundmass is comprised of quartz and subordinate feldspar, biotite, muscovite and pyrite. Relict flow layering is indicated by biotite and muscovite trains which wrap around the phenocrysts (Fig. 6).

Near the top of the volcanics are several thin bands of tremolitic marble consisting of acicular tremolite crystals (up to 10 mm) in a calcite matrix.

Silicate analyses of two rhyolites and one basalt are shown in Table 2. The two rhyolites differ from the world average (column 4) in an enrichment of  $K_2O$  and a deficiency of  $Na_2O$ ,  $Al_2O_3$  and  $CaO$ . Similarly the basalt is enriched in  $K_2O$  and  $Na_2O$  and deficient in  $CaO$  and  $MgO$  compared with the world average. The silica percentage of basalt in column 5 is anomalously high due to numerous quartz-filled amygdales.

(c) Unnamed schists Bd2

Schists outcrop southwest of Peake ruins and north of Mount Denison. They are poorly exposed and outcrop only in creeks. Rock types are pale grey quartz-chlorite-muscovite and quartz-chlorite-epidote schists and phyllites derived by the metamorphism of silty clays. Thin metasilstone and metasandstone lenses, often with graded bedding and cross bedding are common. The beds in the unit have gradational, conformable contacts with the underlying Tidnamurkuna Volcanics and overlying Baltucoodna Quartzite.

(d) Baltucoodna Quartzite Edb

South of Peake ruins and near Mount Denison is a thick sequence of greyish-white quartzites, metabasalts, schists and phyllites.

These rocks, and quartzites, schists, gneisses and calc-silicates occurring in various basement blocks in the Margaret Inlier comprise the Baltucoodna Quartzite (Encl. 1). It has a gradational boundary with the underlying unnamed schist unit and the top is not exposed.

The stratigraphic sequence near Mount Denison has a minimum thickness of 4 500 m. The quartzites are greyish-white and are composed dominantly ( 90%) of quartz. Bedding is defined by numerous metamorphosed clay laminations which range in thickness from 1 - 10 mm. Interbedded pelitic sediments (quartz-muscovite-chlorite schists) rarely exceed 20 m thick.

Interlayered with the quartzites are several basaltic flows and associated sediments (Encl. 1). The basalts are strongly cleaved, fine to medium-grained, and consist of plagioclase, hornblende, epidote, biotite and magnetite. Plagioclase phenocrysts are abundant and vary in size from 2 to 25 mm (Fig. 7). Elongate vesicles and quartz-epidote amygdales are common. The one basalt sample analysed (Table 2, column 6) is similar to the average basalt quoted by Le Maitre (1976). Interlayered with the basalts are dark grey, chloritic phyllites, epidote-actinolite quartzites and rare, thin cream to pale grey limestone.

Quartzites, schists, gneisses and calc-silicates exposed in areas of basement rocks in the Margaret Inlier have tentatively been included within the Baltucoodna Quartzite. In the Coominaree Mine Block quartz-feldspar-biotite and sillimanite gneisses, quartz-biotite schists, plagioclase-hornblende-microcline calc-silicates and a tremolite, talc marble are exposed. A granulite, containing 30% clinopyroxene, outcrops in the War Loan Mine Block.

TABLE 2  
RHYOLITES BASALTS

	1	2	3	4	5	6	7
SiO <sub>2</sub>	74.28	73.48	74.80	72.82	57.19	49.14	49.20
TiO <sub>2</sub>	.32	.31	.46	.28	1.19	1.86	1.84
Al <sub>2</sub> O <sub>3</sub>	12.61	12.29	13.05	13.27	15.41	15.53	15.74
Fe <sub>2</sub> O <sub>3</sub>	1.32	3.00	.99	1.48	5.45	3.91	3.79
FeO	.90	.60	1.56	1.11	4.50	7.20	7.13
MnO	.01	.02	.01	.06	.04	.08	.20
MgO	.46	.28	1.45	.39	3.58	5.85	6.73
CaO	.38	.34	.22	1.14	3.58	8.98	9.47
Na <sub>2</sub> O	1.64	2.58	4.77	3.55	3.65	3.20	2.91
K <sub>2</sub> O	7.24	6.61	1.89	4.30	2.72	1.60	1.10
P <sub>2</sub> O <sub>5</sub>	.08	.03	.05	.07	.13	.42	.35
H <sub>2</sub> O <sup>+</sup>	.57	.27	.99	1.10	1.55	1.63	.95
H <sub>2</sub> O <sup>-</sup>	.13	.13	.60	.31	.17	.21	.43
TOTAL	99.95	99.95	100.84	99.88	99.63	99.60	99.84

1. Rhyolite (Tidnamurkuna Volcanics) near Peake ruins.  
Specimen 6041 RS 145.
2. Rhyolite (Tidnamurkuna Volcanics) near Peake ruins. Specimen  
6041 RS 144.
3. Average of 2 rhyolite analyses. Pepegoona Porphyry (Mount  
Painter Province) A 3830/62 and A 668/65 (Coats & Blissett, 1971).
4. Average of 554 rhyolites (Le Maitre, 1976).
5. Amygdaloidal basalt (Tidnamurkuna Volcanics) near Mount  
Denison. Specimen 6041 RS 146.
6. Basalt (Baltucoodna Quartzite) near Mount Denison. Specimen  
6041 RS 143.
7. Average of 3156 basalts (Le Maitre, 1976).

(e) Unnamed metamorphics Bd3

To the east of the Margaret Inlier are four, small, isolated basement inliers (Encl. 1). Rock types of each inlier differ and include a variety of magnetite-rich psammites, calc-silicates and amphibolites.

Epidote quartzites have the epidote concentrated in thin layers and ellipsoids. The quartzites are whitish-grey, frequently cross-bedded and heavy mineral laminated (Fig. 8). They are also often fractured and brecciated with a plagioclase-amphibole, epidote-magnetite matrix. A quartz-plagioclase-clinopyroxene granulite outcrops at Lagoon Hill. Quartz, quartz and muscovite, and biotite-rich metasandstones and metasiltstones are exposed at Mount Charles and Spring Hill.

Amphibolites present may represent both metavolcanics and metamorphosed calcic sediments. Quartz and quartz-epidote amygdales in various units near Spring Hill suggest a high proportion of metabasalts.

Correlation

The Peake Metamorphics have one possible correlative in South Australia. The Radium Creek Metamorphics in the Mount Painter Province, 350 km to the southeast of the ranges, are dominantly a sequence of arenaceous sediments but also include a possible extrusive acid porphyry (Pepegoona Porphyry: Coats & Blissett, 1971). Silicate analyses of the Pepegoona Porphyry are shown in Table 2 and indicate dissimilar oxide contents to the rhyolites of the Tidnamurkuna Volcanics. No basalts have been observed in the Radium Creek Metamorphics. The correlation between metamorphics of the two terrains must be considered to be very tenuous.



However, the interfingering of volcanics and quartzites so characteristic of all units in the Peake Metamorphics is very similar to the Eastern Creek Volcanics of the Haslingden Group, 800 km to the northeast in the Mount Isa Trough. The Eastern Creek Volcanics, dominantly of metabasalts (often amygdaloidal) and quartzites, are described in detail in Derrick et al., (1976) and Derrick et al., (1977), and are considered to be possibly equivalent to the Peake Metamorphics.

WIRRIECURRIE GRANITE EYw (CARPENTARIAN)

The Wirriecurrie Granite outcrops in the vicinity of Peake Creek. The name 'Wirriecurrie' is derived from the aboriginal word for Peake Creek. The southern and eastern boundaries of the granite are sheared while the northwestern margin is gradational into a zone of migmatites, pegmatites, gneisses and diorites.

The Wirriecurrie Granite consists of gneissic augen granites, porphyritic granites, adamellites, granodiorites and minor aplites. Phenocrysts are ovoid to lenticular bluish quartz (3mm) and rounded to ovoid red potash feldspar and minor plagioclase (30 mm) (Fig. 9). Plagioclase crystals are heavily altered to epidote and sericite while the potash feldspar crystals (which are dominantly perthitic orthoclase and microcline) are only slightly altered (Stevenson, 1973; Webb & Lowder, 1971). Within granites which are strongly foliated, mobilisation of silica has resulted in monomineralic aggregates of quartz exhibiting a granoblastic texture. The foliation is defined by a preferred orientation of biotite. Biotite, epidote and chlorite are only present in minor amounts.

Silicate analyses of two specimens reveal that the

Wirriecurrie Granite is slightly deficient in  $\text{Na}_2\text{O}$  and  $\text{Al}_2\text{O}_3$  and enriched with respect to  $\text{K}_2\text{O}$  relative to world averages for granites and adamellites (Table 3).

TABLE 3  
WIRRIECURRIE GRANITE

	1	2	3	4
$\text{SiO}_2$	70.80	68.65	69.31	71.30
$\text{TiO}_2$	.46	.54	.60	.31
$\text{Al}_2\text{O}_3$	13.50	14.55	13.55	14.32
$\text{Fe}_2\text{O}_3$	.99	1.23	1.15	1.21
$\text{FeO}$	2.90	2.70	3.65	1.64
$\text{MnO}$	.03	.08	.04	.05
$\text{MgO}$	.66	1.14	.78	.71
$\text{CaO}$	1.57	2.68	1.95	1.84
$\text{Na}_2\text{O}$	2.45	3.47	2.58	3.68
$\text{K}_2\text{O}$	5.54	4.00	5.26	4.07
$\text{P}_2\text{O}_5$	.14	.19	.15	.12
$\text{H}_2\text{O}^+$	.77	.59	.95	.64
$\text{H}_2\text{O}^-$	.15	.14	.12	.13
TOTAL	99.95	99.96	100.08	100.02

1. Adamellite (Wirriecurrie Granite) north of Mount Denison.  
Specimen 6041 RS 12
2. Average of 27 adamellites (Le Maitre, 1976).
3. Porphyritic granite (Wirriecurrie Granite) near Peake Creek.  
Specimen 6041 RS 62
4. Average of 197 granites (Le Maitre, 1976).

Diorite dykes form a semicircular pattern in the centre of the granite and have also intruded along faults and the foliation. They vary in thickness from 2m to several hundred metres. Plagioclase and hornblende are the dominant mineral constituents with lesser amounts of quartz, chlorite, epidote and iron oxides. Plagioclase phenocrysts occasionally exhibit a subvariolitic texture.

Geochronology on the Wirriecurrie Granite incorporates nine K-Ar age determinations on biotite and twelve Rb-Sr determinations on total rock samples (Webb & Lowder, 1972; Steveson, 1973; Webb, 1974). The ages can be interpreted as granite intrusion around 1600 Ma overprinted by tectonism during the Musgravian Orogeny (circa 1050 Ma) and Delamerian Orogeny (circa 500 Ma).

A probable correlative of the Wirriecurrie Granite is the Engenina Adamellite in the Mount Woods Inlier 150 Km to the southwest. The Engenina Adamellite is also foliated and porphyritic (Benbow & Flint, 1978) and has been dated by Rb-Sr (total rock) at  $1641 \pm 38$  Ma (Webb, 1977).

#### CALLANNA BEDS (WILLOURAN)

Mawson (1927) originally assigned the name Willouran Series to a sequence in the Willouran Ranges which was very generally described as quartzites, slates and calcareous rocks. Sprigg (1949) adopted the same name to describe the sequence under the Witchelina Quartzite and later proposed that the Willouran Series be included in the Adelaidean System (Sprigg, 1952).

The Callanna Beds as defined by Thomson and Coats (1964) is the name applied to the type Willouran sequence including the underlying sedimentary-volcanic succession now occurring as dismembered blocks in diapiric context. The Callanna Beds are considered to represent the earliest depositional cycle of the Adelaidean System and in the Peake and Denison Ranges are defined by

a basal sequence overlying the Peake Metamorphics with angular unconformity. Formations ascribed to the Callanna Beds are Younghusband Conglomerate (new name), Coominaree Dolomite (new name) and Cadlareena Volcanics (new name) - (see Table 4). The maximum total thickness is at least 800 m.

TABLE 4

Stratigraphy - CALLANNA BEDS

Ecc	Cadlareena Volcanics
Eck	Coominaree Dolomite
Eco	Younghusband Conglomerate

Peake Metamorphics

- conformable contact
- angular unconformity
- ?————— contact uncertain-unknown sequences suspected to intervene.

It is important to note that in the Peake and Denison Ranges disruption always occurs above the Cadlareena Volcanics. As a consequence the position of the upper boundary of the Callanna Beds with the Burra Group is unresolved. Work by Preiss (1974) in the vicinity of the Spalding Inlier near Adelaide (Fig. 3) established that the base of the Burra Group is marked by a heavy mineral laminated sandstone (Rhynie Sandstone), which is in turn overlain by the River Wakefield Subgroup and Bungaree Quartzite (Forbes, 1977). The Mount Margaret Quartzite is tentatively correlated with the Bungaree Quartzite and the Witchelina Quartzite of the Willouran Ranges, but as the equivalent of the Rhynie Sandstone has not been recognised definitely in the Peake and Denison Ranges the position of the base of the Burra Group (and upper boundary of the Callanna Beds) remains uncertain.

As discussed by Thomson (1966) and Preiss (1973), the lower Willouran sequence from the Peake and Denison Ranges closely resembles those of three other regions in which basement rocks are exposed:

- 1) the Mount Painter Province (Coats and Blissett, 1971)
- 2) the Willouran Ranges (Murrell, 1977)
- and 3) Broken Hill area - the Barrier Ranges (Cooper and Tuckwell, 1971).

Table 5 depicts equivalent Adelaidean sequences from the Mount Painter and Broken Hill regions where the basal Willouran sequences correlate closely with those in the Peake and Denison Ranges. A sequence composed of a basal quartzite conglomerate overlain by a dolomite - chert unit in turn overlain by altered amygdaloidal basalts is common to the three regions.

(a) Younghusband Conglomerate Eco

Thomson (1966) first recognised a basal Adelaidean (Willouran) succession unconformably overlying older basement rocks of the Peake Metamorphics about 3 km south of War Loan Mine (War Loan Mine Block). Subsequent mapping by Coats and Preiss in 1972 confirmed the presence of this unconformity and the same Willouran sequence was observed intact, unconformably overlying a small basement inlier 1.2 km north of Coominaree Mine (Coominaree Mine Block). Descriptions of these successions are in Preiss (1973). The lowermost unit in the Willouran succession is herein named the Younghusband Conglomerate (Appendix A, Unit 4). The thickest development and best exposure occurs in the type section on the eastern margin of the Coominaree Mine Block where 27 m of basal clastics are recorded. A detailed description of this section is in Appendix B, Section 1.

The Younghusband Conglomerate consists of two unnamed members. The lower member (12 m thick) consists of a basal sedimentary breccia (2 m thick) overlain by gritty and pebbly sandstones, minor siltstone, coarse dolomitic sandstones and minor red-maroon shales towards the top of the unit. The upper member consists of 15 m of red-brown shales with fine sandstone intercalations. To the south, on the eastern flank of the War Loan Mine Block, a sequence of purple and green-weathering siltstones overlies a basal quartzitic conglomerate-breccia.

Clasts in the basal sedimentary breccia are predominantly vein quartz with minor granite and quartzite pebbles set in a brown feldspathic quartzite matrix. The lithology and high angularity of the clasts suggest a local basement source. Cross-bedding in both members of the Younghusband Conglomerate suggests a shallow water environment of deposition.

To the north on Warrina, about 3.5 km southwest of "Peake" ruins, the southern margin of the Peake Metamorphics is bounded in part by the Younghusband Conglomerate. Only the lower member is present and it outcrops sporadically along a total strike length of 7 km. Graded-bedding and cross-bedding are common. Clast lithologies correspond to local basement rocks, i.e. quartzites, schist, vein quartz, granites and pegmatites with quartzites predominating. A laminated, pebbly, red-brown shale is occasionally interbedded with the conglomerate. Outcrop is often in the form of isolated residual cappings separated from and lying topographically above the Cadlareena Volcanics (Fig. 10). These cappings, resting unconformably on Peake Metamorphics, represent erosional remnants and provide evidence of a minor pause in sedimentation between the Younghusband Conglomerate and Cadlareena Volcanics. The absence of the Coominaree Dolomite,

at least to the northwest, confirms the presence of an hiatus above the basal conglomerate in this northern part of the ranges. Drilling by Uranerz (Iliff, 1975) intersected 7 m of basal clastics on the northwestern margin of the conglomerate outcrop which is the maximum thickness recorded on Warrina.

(b) Coominaree Dolomite Eck

The Younghusband Conglomerate is conformably overlain by a sequence of mainly stromatolitic dolomite herein named the Coominaree Dolomite (Appendix A, Unit 5). The most complete succession (thickness of 77 m) occurs 1.1 km north-northeast of Coominaree Mine where a detailed section was described by Preiss, (1973). This type section is described in Appendix B, Section 2.

The lower part of the Coominaree Dolomite is predominantly buff and pale brown dolomites with gritty and pebbly layers displaying low angle cross-bedding. Overlying this is a series of flaggy, buff dolomites and fine sandstones which pass upwards into stromatolitic dolomites, then non-stromatolitic dolomites, oolitic at the base, and finally, an upper stromatolitic dolomite. The upper part of the formation is poorly exposed although stromatolitic float is observed. The stromatolites Acaciella cf. australica and Gymnosolen cf. ramsayi occur in this formation and are described by Preiss (1973).

On the eastern flank of the War Loan Mine Block the Coominaree Dolomite conformably overlies the Younghusband Conglomerate and passes upwards into the Cadlareena Volcanics. This sequence was first studied by Thomson (1966) and later by Coats and Preiss (1973). To the north on Warrina the Coominaree Dolomite is largely absent, being present only on the southeastern margin of Younghusband Conglomerate outcrop.

Here a thin bed (2 m thick) of pebbly, silty dolomites outcrops along a strike length of 100-150 m before lensing out to the northwest. Stromatolitic dolomites, typical of the Coominaree Dolomite, also occur in a small rafted block in a diapir about 1 km to the west of this outcrop.

The Coominaree Dolomite probably was deposited in a shallow water/shelf environment but the thin pebbly dolomite in places underlying volcanics may represent a tidal channel facies deposited on the margin of a shallow basin. Along strike to the northwest, Cadlareena Volcanics disconformably overlie Younghusband Conglomerate and the Coominaree Dolomite is absent.

(c) Cadlareena Volcanics Ecc

Amygdaloidal lavas of Willouran age were first recognised in the Peake and Denison Ranges on Anna Creek, near Douglas Well, by Wopfner and Heath in 1962. Similar rocks were described by Thomson (1966) in a basal Adelaidean sequence outcropping northeast and southeast of Nilpinna H.S. and also about 45 km to the north on Warrina. This volcanic sequence is herein named the Cadlareena Volcanics (Appendix A, Unit 6), a name first used informally by H. Wopfner. The lower part of the type section is on Warrina where a basal contact with the Younghusband Conglomerate is preserved. The upper part of the type section is best recorded on Anna Creek near Douglas Well. Detailed descriptions of both these sections are in Appendix B, Sections 3A, 3B.

In addition to the outcrops in stratigraphic context, a number of rafted blocks of volcanics (varying in size from a few to several hundred metres across) occur in diapirs, most commonly in the northern part of the ranges.



A typical example is along the eastern margin of the Rockwater Hill Block where a number of rafted volcanic blocks (too small to be shown on the map) occur within a diapir but are aligned parallel to the basement - diapir contact. As a result of diapirism associated with decollement at the base of the Adelaidean, these scattered volcanic remnants provide the sole evidence of an original basal Adelaidean sequence at this particular locality.

The volcanics thin northwards, from a maximum thickness of 730 m at Douglas Well to approximately 200 m on Warrina. It should be emphasized that throughout the ranges diapirism occurs above the volcanics and hence these thicknesses are minimum estimates considering the uncertainty of the upper contact. The volcanics extend northwards at least as far as Mount Dutton where a small block of volcanics outcrops in a fault zone.

Altered basaltic to andesitic volcanics tentatively correlated with the Cadlareena Volcanics were intersected in Boorthanna No. 1, 30 km southwest of the ranges (Fig. 3) (Holmes, 1970; Holmes and Rayment, 1970).

On Warrina the volcanics generally disconformably overlie the Younghusband Conglomerate (Fig. 10) and more rarely basalts lie directly on rocks of the Peake Metamorphics (Fig. 11). On the southern part of Warrina and on Boorthanna the volcanics conformably overlie the Coominaree Dolomite.

The Cadlareena Volcanics are a complex series of mainly basaltic lava flows with individual flows 20-30 m thick, and thin interbedded sediments and pyroclastics (Fig. 12). Associated with the basalts are dolerites and andesites plus rare thin rhyolites and dacites. Several volcanic rocks of trachytic composition have been identified and are thought

to be products of potash metasomatism (Radke, 1976; 1977). Whitehead (1972b) recognised secondary microcline in basalts from Douglas Well. The presence of tourmaline in several samples indicates boron metasomatism.

The lavas are predominantly amygdaloidal. The abundance and size of the amygdales are variable within individual flows and are generally at a maximum at the tops of flows. Plagioclase (often albite), orthoclase, chlorite, zeolites, tremolite-actinolite, epidote, calcite, quartz and fine-grained basalt infill the amygdales. Whitehead (1972) showed amygdales to contain varying proportions of these minerals with the order of crystallisation as follows: albite-epidote-calcite-chlorite-quartz. The groundmass of the basalts generally consists of randomly orientated plagioclase laths together with abundant amphibole and chlorite (Radke, 1976).

All of the volcanics have suffered extensive alteration (mostly deuteric) although the original igneous textures are generally well preserved. Alteration processes include:-

- 1) alteration of calcic plagioclase to albite-andesine,
- 2) muscovite-sericite replacing plagioclase,
- 3) fibrous tremolite-actinolite replacing pyroxene,
- 4) chlorite and opaque minerals replacing amphibole, pyroxene and ?olivine,
- 5) secondary microcline replacing plagioclase.

Interstitial granular and microcrystalline silica has been introduced during weathering.

The volcanics on Warrina have not been previously studied in detail. They disconformably overlie the Younghusband Conglomerate and outcrop about 4 km southwest of "Peake" ruins along a northwest strike length of 9 km. Thickness is about

200 m in the southwest, decreasing to less than 50 m along strike to the northwest. Thin, lenticular mudstones are interbedded with basalts at or near the base of the unit (Appendix B, section 3A). The mudstones are usually red-maroon and are occasionally sandy and more rarely pebbly. The volcanics are mainly dark green-grey amygdaloidal basalts, dolerites and andesites with interbedded lapilli, and maroon tuffs and tuffaceous sediments. The proportion of pyroclastics and tuffaceous sediments generally increases towards the top of the unit as indicated by the drilling results of Uranerz (Iliff, 1975) which recorded a thin volcanic breccia near the top of the sequence.

Several basaltic flows were identified, each displaying the following sequence of rock types (C.D. Branch, pers. comm.):

Top 1) coarsely vesicular basalt

2) coarse-grained non-vesicular dolerite

Base 3) lapilli tuffs and intertonguing tuffaceous sediments.

Estimated thickness of the flows is 20-30 m but their exact thickness and number were not measured due to rapid horizontal and vertical variability.

The upper part of the volcanics is often marked by a thin, lenticular, rippled, red-brown mudstone or quartzite. Volcanic clasts in the mudstones indicate reworking of the basalts in a shallow water environment.

The volcanic sequence at Douglas Well on Anna Creek (Appendix B, Section 3B) is similar to that on Warrina but important differences and characteristics of the Cadlareena Volcanics at Douglas Well are:

- a) the sequence is at least 730 m thick. The base and top of the sequence are not exposed;

- b) a 5-10 m thick medium-coarse sandstone with numerous clay pebbles occurs low in the sequence;
- c) up to 100 m of quartzite and sandstone (some coarse-grained) marks the upper part of the formation;
- d) several thin, lenticular rhyolites outcrop near the top of the sequence;
- e) secondary silicification is extensive.

On Boorthanna, in the vicinity of Warloan Mine, at least 300 m of volcanics outcrop in a synclinal structure. Thin rhyolites have been recognised in the mainly basaltic sequence.

Immediately northeast of Coominaree Mine on Boorthanna the basal Adelaidean sequence is preserved lying unconformably on Peake Metamorphics (Coominaree Mine Block) and includes several hundred metres of volcanics conformably overlying the Coominaree Dolomite.

The volcanics near Douglas Well were first correlated with the Woollana Volcanics of the Mount Painter Province by H. Wopfner (Wopfner in Thomson, 1966). In both areas the basal Adelaidean stratigraphy can be generalised into a basal conglomerate and an overlying dolomite unit which passes into volcanics. The similarity of stratigraphic position and mineralogy of the volcanics in the two regions is the basis for the correlation. Other aspects common to both formations are:

- a) the volcanics consist of a complex series of flows with minor interbedded sediments;
- b) interbedded sediments include lapilli tuffs, tuffs, volcanic breccia (minor) plus shallow water sandstones and mudstones;
- c) amygdaloidal lavas with associated basaltic textures (randomly orientated plagioclase laths);

- d) original mineralogy of pyroxene, hornblende and olivine has been extensively altered to tremolite-actinolite, epidote, chlorite, calcite and opaque minerals;
- e) potash metasomatism and albitization are evident to varying degrees;
- f) the volcanics are iron rich and show secondary enrichment of iron oxides plus late stage silicification;
- g) similarity of silicate and trace element analyses (Tables 6 & 7);
- h) minor acid volcanics (rhyolite) occur in the mainly basaltic sequences.

Crawford (1963) considered the Wooltana Volcanics to be sub-aerially extruded from fissures with explosive activity from central vents supplying the pyroclastic material. A similar environment is envisaged for the Cadlareena Volcanics but with explosive activity and shallow water sedimentation increasing towards the top of the unit.

On Warrina minor copper and uranium mineralisation has been identified in the Cadlareena Volcanics. Aerial surveys by Uranerz first revealed radiometric anomalies and these were followed up by extensive carborne and footborne scintillometer surveys along the unconformity between the volcanics and Peake Metamorphics (Iliff, 1975). The base of the unit showed several radiometric anomalies. Samples taken from trenches on the western margin of the volcanics assayed up to 100 ppm uranium, and minor malachite and azurite staining was observed in one of the trenches. Drilling, however, yielded no significant mineralisation. Copper carbonates were observed in one locality in the basal mudstone and faulting nearby suggests a hydrothermal origin for the mineralisation. Trace element contents

of volcanics from Warrina and Anna Creek (Douglas Well) are listed in Table 7. Uranium contents are uniformly low as are the mean base metal contents which are generally less than those quoted for an average basaltic rock by Turekian and Wedepohl (1961).

A summary of the major element chemistry of the Cadlareena Volcanics and the Wooltana Volcanics (from the Mount Painter Province and Depot Creek) is presented in Table 6. The average composition of continental basalts (column 6) applies to basalts erupted through continental crust at continental rifts (Pearce, 1976).

TABLE 6

	1.	2.	3.	4.	5.	6.	7.
SiO <sub>2</sub>	48.48	47.71	48.43	48.22	48.54	48.81	66.00
Al <sub>2</sub> O <sub>3</sub>	13.05	12.97	13.08	13.85	13.79	14.41	14.22
Fe <sub>2</sub> O <sub>3</sub>	10.64	9.63	9.08	12.33	13.29	12.00	3.53
FeO	3.84	1.88	3.74	2.91	-		.35
MnO	0.12	0.09	0.10	0.25	.04	5.96	.01
MgO	5.86	8.25	6.92	7.26	9.21	10.05	.29
CaO	6.94	8.24	8.86	4.71	0.36	2.90	.87
Na <sub>2</sub> O	3.38	3.35	3.04	1.80	0.21	-	.23
K <sub>2</sub> O	1.97	1.65	1.66	3.66	5.61	0.95	12.29
P <sub>2</sub> O <sub>5</sub>	0.16	0.13	0.15	0.13	0.14		.18
TiO <sub>2</sub>	1.99	1.46	1.63	1.55	1.76	2.47	.70
H <sub>2</sub> O <sup>+</sup>	2.37	2.59	2.57	3.03	-	-	.44
H <sub>2</sub> O <sup>-</sup>	0.34	0.28	0.24	0.25			.16
CO <sub>2</sub>	0.85	-	0.05	0.67			
S	0.04						
TOTAL	100.03	98.21	99.55	100.62	92.95	97.55	99.27

1. Cadlareena Volcanics from Anna Creek (Douglas Well). Average of 7 analyses; 6140 R.S. 4-8, 23 and 25. Range SiO<sub>2</sub> 45.1-49.4, Al<sub>2</sub>O<sub>3</sub> 11.96-13.8, total Fe 12.25-17.1, MgO 2.7-7.63, CaO, 5.2-9.44, Na<sub>2</sub>O 2.94-4.7, K<sub>2</sub>O 0.33-4.7
2. Cadlareena Volcanics from Warrina. Average of 4 analyses;

6041 R.S. 161-164. Range  $\text{SiO}_2$  46.57-49.84,  $\text{Al}_2\text{O}_3$  11.93-14.67, Total Fe 10.3-12.8, MgO 11.53-5.24, CaO 5.1-11.99,  $\text{Na}_2\text{O}$  1.7-4.45,  $\text{K}_2\text{O}$  1.13-2.41.

3. Average of Cadlareena Volcanics from Warrina and Anna Creek (as above) plus 6040 R.S. 14 and 15 from Boorthanna. Average of 13 analyses. Range  $\text{SiO}_2$  45.1-49.84,  $\text{Al}_2\text{O}_3$  11.93-14.67, total Fe 10.3-17.1, MgO 2.7-11.53, CaO 5.1-15.4,  $\text{Na}_2\text{O}$  0.48-4.7,  $\text{K}_2\text{O}$  0.33-4.7.
4. Wooltana Volcanics from the Mount Painter Province-average of 15 analyses (Coats, in prep). Range  $\text{SiO}_2$  37.8-59.2,  $\text{Al}_2\text{O}_3$  10.8-18.76, total Fe 8.25-37.0, MgO 0.22-12.68, CaO 0.04-15.8,  $\text{Na}_2\text{O}$  0.13-4.6,  $\text{K}_2\text{O}$  0.15-10.2.
5. Wooltana Volcanics from Depot Creek. Average of 8 analyses, 6433 R.S. 2,3,5,6,7,8,9 and 10. Range  $\text{SiO}_2$  40.5-55.1,  $\text{Al}_2\text{O}_3$  10.4-20.2, total Fe 2.6-20.3, MgO 3.2-11.2, CaO 0.16-0.57,  $\text{Na}_2\text{O}$  0.05-0.72,  $\text{K}_2\text{O}$  2.5-11.0.
6. Average distribution of elements in a continental basalt (Pearce, 1977).
7. Rhyolite from Cadlareena Volcanics on Anna Creek; 6140 R.S. 24.  
Average analyses of Cadlareena Volcanics from Warrina and Anna Creek (Table 6, columns 1 & 2) are similar. The average compositions of the Cadlareena and Wooltana Volcanics are similar (Table 6, columns 3,4 & 5), although there are variations in the alkali contents. The Cadlareena Volcanics are less potassic and more calcic than the Wooltana Volcanics and the rocks from Depot Creek are particularly low in Ca and Na and correspondingly high in K. As noted by Thomson (1966) the Cadlareena Volcanics are characterised by a MgO/CaO ratio of less than one while MgO/CaO ratios for the Wooltana Volcanics exceed one.

Fander (1963) noted the highly variable alkali content of the Wooltana Volcanics and generally described the lavas as sodic trachytes with subordinate andesites and rhyolites. Turner (1969) summarized the Wooltana Volcanics as belonging to the trachyte family with a late-stage mineral assemblage which is predominantly potash-rich. This contradiction in itself testifies to the highly variable alkali content of these volcanics which is largely due to deuteric alteration (e.g. albitization) and potash metasomatism.

Whitehead (1972b) described the Cadlareena Volcanics as spilites (Na enriched), while Radke (1976) used the term 'altered basalts'. Both writers recognised minor potassium-rich varieties and considered them to be products of potash metasomatism and this, together with the type of deuteric alteration, are features common to both the Cadlareena and Wooltana Volcanics.

Plotting analyses of Cadlareena and Wooltana Volcanics on an AFM diagram (Fig. 13) does not reveal any differentiation trends. Most of the analyses plot in a restricted zone but are scattered in an irregular manner. This scatter may reflect alteration of the volcanics which effectively masks any pre-existing differentiation trends. One can generally conclude that the major element compositions of the Cadlareena and Wooltana Volcanics are broadly analogous and notable variations in Na, K and Ca abundances occur within both units.

A summary of the trace element geochemistry of the Cadlareena and Wooltana Volcanics is presented in Table 7. Analyses of volcanics from Warrina and Anna Creek (Douglas Well) are broadly analogous (Table 7,) although the rocks from Douglas Well appear enriched in Ba. The Cadlareena Volcanics as a whole are depleted



in Cu and Zn relative to "average" basaltic rocks (Turekian and Wedepohl, 1961) but Ni, Pb, Co, Li, Ba, Zn, U and Th values generally correspond (Table 7, lines 3 & 5). The highest Cu (80 ppm) and Pb (800 ppm) values occurred in a dacite from Warrina (6041 RS 93). Uranium values are uniformly low although several andesites on Warrina recorded uranium contents up to 30 ppm (6041 RS 100).

The Wooltana Volcanics have higher average Cu, Pb and Zn contents than the Cadlareena Volcanics while Rb and Sr contents are similar.

TABLE 7

	Zn	Cr	Mo	Cu	Ni	Co	Li	Pb	Ag	Rb	Sr	V	Ba	Zr	U	Th
1.	27	199	8	18	197	39	40	81	-	55	162	251	202	87	6	<4
2.	51	375	8	21	93	37	25	27	0.2	116	114	600	1448	113	<4	4
3.	43	335	6	20	125	40	29	19	0.2	92	142	407	416	99	<4	<4
4.	183			31	155	57		8	0.2			128	156			
5.	105	170	1.5	87	130	48	17	6	0.1	30	465	250	330	140	1	4

1. Cadlareena Volcanics (6041 R.S. 93, 95-101, 161-164). Mean of 12 samples from Warrina.
2. Cadlareena Volcanics (6140 R.S. 4-8 and 23-28). Mean of 8 samples from Anna Creek (Douglas Well).
3. Cadlareena Volcanics-mean of all analyses from Warrina, Anna Creek and Boorthanna (6040 R.S. 14-15).
4. Wooltana Volcanics-mean of 8 analyses (Coats & Blissett, 1971).
5. Distribution of elements in basaltic rocks (Turekian & Wedepohl, 1961).

#### SEQUENCES OF UNCERTAIN AGE (WILLOURAN OR TORRENSIAN)

Block faulting and diapirism have caused dislocation and disruption of most of the Adelaidean sequences above the Cadlareena Volcanics. No complete succession is observed from the Callanna Beds upwards into the Burra Group. At least 16 000 m of sediments intervene between the youngest recognised unit in the Callanna Beds (Cadlareena Volcanics) and the oldest recognised unit in the Burra Group (unnamed siltstone). Reconstruction of the stratigraphy of

this considerable thickness of sediments is a difficult proposition due to the structural contexts in which most of the rocks occur, i.e. in rafted blocks in diapirs and in faulted blocks. Since none of these rocks can be ascribed with certainty to either the Callanna Beds or the Burra Group they are described under the heading "Sequences of Uncertain Age". The units involved are the Rockwater Beds, War Loan Beds, Nilpinna Beds, Duff Creek Beds and Murrana Beds (Table 8).

TABLE 8

Stratigraphy - SEQUENCES OF UNCERTAIN AGE

Ea	Undifferentiated blocks
	_____?_____?
Eaa	Murrana Beds
	-----?-----?-----
Eaf	Duff Creek Beds
	-----?-----?-----
Ean	Nilpinna Beds
	-----?-----?-----
Eaw	War Loan Beds
	-----?-----?-----
Ead	Rockwater Beds
	_____?_____?

CALLANNA BEDS - Cadlareena Volcanics

\_\_\_\_\_?\_\_\_\_\_?

contact uncertain: unknown  
sequences suspected to intervene

-----?-----?-----?---

transition suspected on basis of  
lithological overlap

The stratigraphic succession is far from complete. Lithological overlaps between spatially separated sequences are the main criteria used in assembling the stratigraphic sequence. Some of the gaps in the succession portrayed in Table 8 may be filled, at least in part, by miscellaneous sequences described under the sub-heading "Undifferentiated Blocks". These rock units are not formally named and occur in thin sequences in blocks in diapirs and in small faulted blocks.

Equivalents of the Rockwater Beds and Duff Creek Beds occur in the Willouran Ranges (R.P. Coats, pers. comm.). It is probable other units described in this section also have equivalents in the Willouran Ranges.

(a) Rockwater Beds Bad

The Rockwater Beds (Appendix A, Unit 7) outcrop in four small blocks, all tectonically isolated in disrupted diapiric zones. Distinctive black cherts (which weather white) and black dolomites characterise the unit. The Rockwater Beds are believed to be equivalent to the Dunn's Mine Beds found in the Willouran Ranges (Murrell, 1977).

Two small blocks outcrop on Warrina in a broad disrupted zone above the Cadlareena Volcanics about 11 km northeast of Warrina R.S. A third block outcrops immediately south of Coominaree Mine.

These blocks are composed of sequences of blue-grey to black cherts, grey-black pebbly dolomites, black shales, siltstones, and quartzitic sandstones. The largest block contains about 100 m of sediments. Similar distinctive lithologies occur in several small blocks outcropping in a disrupted diapiric zone on the western margin of the Rockwater Hill Block. Small outcrops of brecciated Cadlareena Volcanics also occur in this zone which probably represents remnants of a lower Adelaidean sequence disrupted by diapirism.

It is noteworthy that all blocks containing Rockwater Beds sequences, although tectonically isolated, outcrop very close to the Cadlareena Volcanics. This association which in itself suggests the Rockwater Beds lie in close proximity stratigraphically to the top of the volcanic sequence, is supported by the fact that the Dunn's Mine Beds in the Willouran Ranges occur in a sequence closely overlying the volcanics (Murrell, 1977).

(b) War Loan Beds     Paw

A sequence of 600 m of shales, sandstones and minor dolomites outcropping in a synclinal structure adjacent to the War Loan Mine, 8 km southeast of Nilpinna H.S., is herein named the War Loan Beds (Appendix A, Unit 8). A detailed description of the type section is in Appendix C, Section 4.

The War Loan Beds exhibit tectonic contacts with the Cadlareena Volcanics in the type section immediately south of War Loan Mine and consequently the lower boundary of the sequence is unresolved. The section headed "Undifferentiated Blocks" describes a total of 5 000 m of sediments which occur in faulted blocks and as rafted blocks in diapirs. It is probable that at least some of these rocks belong to sequences intervening between the Rockwater Beds and War Loan Beds.

The lower part of the War Loan Beds consists of laminated bluish-grey shales, buff weathering dark grey dolomites and minor cross laminated flaggy siltstones. Green-grey and blue-grey shales, brown friable sandstones and greenish gritty feldspathic sandstones characterise the upper part of the section. These lithologies have many affinities with the lower part of the Nilpinna Beds although the only contact observed between the two units (1km north of War Loan Mine) is sheared. Despite the tectonic nature of this contact, overlap of lithologies suggests a passage from the War Loan Beds into the Nilpinna Beds (Table 8).

(c) Nilpinna Beds     Pan

A thick sequence of laminated sandstones and silty shales, outcropping in a fault-bounded sequence 1 km north of War Loan Mine, is herein named the Nilpinna Beds (Appendix A, Unit 9). This unit probably intervenes between the War Loan Beds and Duff Creek Beds.

The type section, 6 km southeast of Nilpinna H.S., exhibits sheared contacts at the base and top with the War Loan Beds and

Duff Creek Beds respectively. The Nilpinna Beds are approximately 2100 m thick and a description of the type section is in Appendix C, section 5. Main lithologies are thinly bedded sandstones (festoon cross-bedding, salt casts - Fig. 14), green arkoses, grey-green silty shales (ripples, mud cracks and salt casts) and minor grey silty dolomites towards the top of the unit.

Platy friable sandstones (salt casts, cross-beds) near the top of the sequence strongly resemble sandstones in the basal Duff Creek Beds. Blue-grey shales and fine-medium sandstones (festoon cross-bedding, salt casts) near the base of the Nilpinna Beds have many affinities with units in the upper War Loan Beds. These lithological criteria are considered to indicate the Nilpinna Beds record an interval of sedimentation between the War Loan Beds and Duff Creek Beds. The transition upwards into the Duff Creek Beds is marked by a gradual increase in the frequency of dolomites.

(d) Duff Creek Beds      Eaf

Reyner (1955) defined the Duff Creek Formation to include all Willouran rocks of the Peake and Denison Ranges. The term Duff Creek Formation has since been abandoned and the name Callanna Beds is used to describe the earliest Adelaidean sediments.

The term Duff Creek Beds was introduced by Thomson et al., (1964) to apply to a unit including green and carbonaceous shales and thin distinctive yellow dolomites. Approximately 3 700 m of shallow water sediments outcrop in the type section on Warrina, east of Nilpinna Homestead. Subsequent mapping revealed that an additional 1 800 m of similar sediments conformably underlie this sequence about 9-10 km to the south on Boorthanna. It is proposed that the type section of Thomson & Coats (1964) be extended to include these sediments, resulting in a total thickness of 5 500 m (Appendix C, Sections 6A & 6B).

This is a minimum thickness since both the upper and lower limits of the type section are bounded by faults.

Lithologies characteristic of the unit include:

1. fine sandstones with platy bedding, ripples, cross-bedding and rare salt casts. Pyritic sandstones occur in the lower half of the section;
2. olive green siltstones-shales with ripples, mud cracks and rare salt-cast horizons. Thin arkosic grits with characteristic blue quartz are often interbedded with these siltstones and shale;
3. flaggy, pale orange-buff weathering, pale grey-yellow dolomites, often gypsiferous, pyritic and occasionally stromatolitic (Fig. 15). Most of the stromatolites are a flat laminated type. The dolomites are occasionally sandy and more rarely conglomeratic and contain intraformational dolomite clasts;
4. medium-coarse gritty sandstone (sometimes feldspathic) and quartzites;
5. black pyritic shales plus occasional brown to buff weathering black dolomites.

Shallow water conditions in lagoonal, supratidal and intertidal environments are indicated by numerous sedimentary structures including mud cracks (Fig. 16), cross-bedding, ripple marks, graded-bedding, flute casts and salt casts. Black, pyritic shales and dolomites were deposited in a reducing lagoonal environment. Pale, gypsiferous dolomites and fine-grained siltstone-sandstone units containing salt and gypsum casts are indicative of a supratidal environment while quartzites and cross-bedded sandstones suggest intertidal shoreline conditions.

The large thickness of shallow water sediments preserved (5500 m) attests to considerable basinal subsidence during deposition. Terrigenous clastic-carbonate couplets indicate cyclic sedimentation in response to sea level fluctuations. Regressive, upward-shoaling cycles (Wilson, 1976) account for at least part

of the large thickness of shallow water sediments preserved.

The dolomites are most likely the result of early diagenetic (syngenetic) dolomitization by seawater. Brines in tidal flats, lagoons and supratidal zones (sabkas), often gain high Mg/Ca ratios due to loss of Ca through evaporative precipitation of gypsum and anhydrite. These Mg-enriched brines migrate down through unconsolidated lime mud dolomitizing the sediment. Extensive dolomitization through thick sections may be aided by the tendency towards prograding sedimentation (Wilson, 1976).

A number of blocks containing rocks belonging to the Duff Creek Beds occur in faulted and diapiric contexts. The largest of these are described below.

- a) Approximately 1 100 m of Duff Creek Beds occur in a faulted block immediately west of Tarlton Springs on Anna Creek. Pyritic green-grey shale, pale yellow-brown gypsiferous dolomite, fine sandstone, quartzite, thin arkose and minor dark grey dolomite represent the main rock types. Flat laminated stromatolites, ripple marks, cross-bedding, mud cracks and minor salt casts are characteristic.
- b) Approximately 1 800 m of sediments belonging to the Duff Creek Beds outcrop in a faulted block about 5.5 km northwest of Douglas Well. Main lithologies include pale stromatolitic dolomites, quartzites, fine sandstones, olive green shales and arkoses. A pebbly quartzite containing fragments of Peake Metamorphics (quartz, quartzite, granite and pegmatite) occurs in the lower part of the section. The lowermost 800 m of the section consists of arenaceous-argillaceous sediments with quartzites, sandstones and shales predominating. Mud

cracks, ripple marks, salt casts and cross-bedding are common.

- c) Approximately 5 km west of Mount Denison on Warrina a number of disconnected blocks occur in a diapiric zone. The sediments are strongly folded, with quartz-biotite schists and slaty argillaceous quartzites predominating, with minor carbonates.
- d) Immediately southeast of Mount Kingston approximately 800 m of green slaty pyritic siltstone, fine sandstone, quartzite and minor pyritic dolomite and arkosic grit occur in an anticline faulted against basement rocks to the west. Similar but smaller blocks are infaulted against basement rocks to the south of "Peake" ruins along the eastern faulted margin of the Denison Inlier.
- e) The southern part of the Dutton Inlier consists of 250 m of Duff Creek Beds. Coarse cross-bedded arkoses, olive green siltstones, fine sandstones and pale stromatolitic dolomites predominate. A laminated gritty quartzitic sandstone at the top of the sequence has many affinities with units in the Murrana Beds.
- e) Murrana Beds Paa

A thick sequence of arenaceous sediments, which probably grade downwards into the Duff Creek Beds, is herein named the Murrana Beds (Appendix A, Unit 11). The type section, about 4 km northwest of Warrina R.S., comprises the eastern limb of a faulted, basinal syncline. The total thickness of the type section is 2 900 m (Appendix C, Section 7).

The lower 700 m of the type section consist of laminated gritty quartzites, fine sandstones, and feldspathic sandstones with grey-green silty shale intercalations.



This lower part of the section has been partly disrupted. Crumpled and dislocated bedding, faulting and minor intrusion of carbonate breccia are characteristic. However, in this lower disrupted zone the general maintenance of bedding orientations and lithological types consistent with the undisturbed overlying sequence indicates stratigraphic continuity.

The lower disrupted zone grades upwards into an undisturbed sequence of mainly arenaceous and silty sediments approximately 1 800 m in thickness. Gritty quartzitic sandstones and interbedded grey-green silty shales are characteristic. Cross-bedding (small and medium scale), ripples, flute casts, mud cracks and graded bedding are ubiquitous. Salt casts are less common.

The uppermost 400 m consists of green nodular siltstones (rarely blue-grey), massive pebbly dolomites, fine to coarse gritty feldspathic sandstones, laminated quartzites, purple and maroon shales and a lenticular, pebbly ?stromatolitic dolomite near the top. The sedimentary structures and lithologies observed in the type section are consistent with deposition in a shallow marine environment.

In the Dutton Inlier a thick sequence of Murrana Beds is separated from Duff Creek Beds by a thin wedge of diapir. A gritty quartzitic sandstone at the top of Duff Creek Beds sequence strongly resembles quartzites characteristic of the Murrana Beds. Similarly, near the base of the Murrana Beds are a few thin beds which are reminiscent of the Duff Creek Beds. Thus although no sedimentary contact is observed between the two units these lithological criteria suggest a passage from the Duff Creek Beds into the Murrana Beds marked by an increased sand-silt:carbonate ratio.

On Warrina both 9.1 km north of Warrina R.S. and 1.2 km east of Nilpinna H.S. the Murrana Beds are observed in tectonic contact with sequences of Duff Creek Beds. Both units have the same facing and orientation and are separated by a thin zone of brecciation.

f) Undifferentiated Blocks    Ea

Rocks from sequences of unknown age, described below, occur in two general contexts:

- 1) in fault-bounded blocks of various sizes;
- 2) in rafted blocks bounded by carbonate breccia occurring in disrupted 'diapiric' zones. One such zone occurs above the Cadlareena Volcanics on Warrina. Sediments from the larger blocks in this zone are described below (see Encl. 1 for block number localities).
  - a) Block 1 consists of approximately 200 m of maroon shales (zoned salt casts), interbedded fine silty sandstones (ripples, cross-beds) and occasional feldspathic sandstones.
  - b) Block 2 comprises sandstones and quartzites interbedded with minor stromatolitic dolomites, purple shales and siltstones.
  - c) Block 3 contains a sequence of pale dolomites and dolomitic siltstones, laminated grey quartzites, green siltstones and rippled sandstones.
  - d) Block 4 consists of mainly cross-laminated quartzites and sandstones with interbedded orange weathering dolomites, grey-green flaggy siltstones and minor maroon shales.
  - e) Block 5 consists of black shales, olive green silty shales and buff dolomites.
  - f) Block 6, at the head of Levi Creek, consists of interbedded coarse feldspathic sandstones, quartzites and grey-green

slaty siltstones (salt casts, ripples and cross-bedding).

- g) Block 7, immediately northeast of Rockwater Hill, consists of quartzites, sandstones, siltstones and minor sandy stromatolitic dolomites.
- h) Block 8 is faulted against basement rocks north of Mount Kingston. The sequence is 700 m thick, the lower 500 m consisting of coarse pebbly arkoses and sandstones overlain by black shales and dolomites (pyritic) with some pale brown dolomites and blue-grey siltstones. The upper 200 m consists of green-grey siltstones (gritty), fine-coarse sandstones and minor cross-laminated quartzites.

The exact stratigraphic positions of the sediments described in blocks 1-8 are unresolved. These rocks only occur in faulted or diapirically rafted blocks and are not observed in thick sequences anywhere in the Peake and Denison Ranges. The total thickness of sediments in these blocks is approximately 5000 m. Obviously a significant part of the lower Adelaidean stratigraphy has been dismembered by diapirism and block faulting. The sediments in the blocks described probably represent remnants of sequences which occur somewhere above the top of the Cadlareena Volcanics (Callanna Beds) and below the recognisable base of the Burra Group (unnamed siltstone). This was concluded after considering:

- 1) the close spatial relationship between Willouran rocks (e.g. Cadlareena Volcanics) and many of the blocks described;
- 2) the fact that diapirism has disrupted lower Adelaidean sequences in other parts of the Adelaide Geosyncline;
- 3) the blocks described above contain no sediments which correspond with upper Torrensian and Sturtian sequences observed in the Peake and Denison Ranges.

# BURRA GROUP (TORRENSIAN)

The following units comprise the Burra Group in the Peake and Denison Ranges; unnamed siltstone, Fountain Spring Beds (new name), Mount Margaret Quartzite, Skillogalee Dolomite, unnamed transition unit and Kalachalpa Formation (new name) (Table 9). The sediments are almost entirely shallow marine and total at least 10 000 m in thickness.

TABLE 9  
Stratigraphy BURRA GROUP

Ebh	Kalachalpa Formation
Eb2	unnamed transition unit
Ebk	Skillogalee Dolomite
Ebo	Mount Margaret Quartzite
Eb1	Fountain Spring Beds
EbI	unnamed siltstone

## SEQUENCES OF UNCERTAIN AGE

- \_\_\_\_\_ conformable contact
- ?--?--?-- transitional contact suspected on basis of lithological overlap
- \_\_\_?\_\_\_?\_\_\_? contact uncertain - unknown sequences suspected to intervene
- ~~~~~ disconformity

As previously discussed, the Rhynie Sandstone, with its characteristic heavy mineral laminations provides a useful marker for the base of the Burra Group in the southern part of the Adelaide Geosyncline (Preiss, 1974). This heavy mineral laminated unit has not been recognised in the Peake

and Denison Ranges and the lowermost unit attributed to the Burra Group (EbI- unnamed siltstone) outcrops in tectonically isolated blocks. Consequently the base of the Burra Group cannot be defined. Units observed in the "unnamed siltstone" (EbI) have lithological affinities with sequences previously described under the heading "Sequences of Uncertain Age" and at least some of these sequences may belong in the lower Burra Group.

Burra Group sediments are restricted in outcrop to the Margaret Inlier where they outcrop in a series of basinal synclines forming the central core of the inlier. The synclines have axial plane traces generally aligned north-northwest which form an en echelon pattern, stepped over to the east as the axes are traced from south to north. Upper Burra Group sediments are restricted in outcrop to folded sequences exposed on the southwest margin of the Margaret Inlier.

(a) Unnamed siltstone Eb1

A sequence of siltstones, fine sandstones and silty dolomites, totalling at least 1 200 m in thickness, comprise the lowermost part of the Burra Group recognised on WARRINA. The sequence has been dismembered by faulting and occurs in several blocks outcropping on the eastern side of the Margaret Inlier, about 4.5 km north of Mount Margaret. The stratigraphy is obscured by a combination of faulting, folding and diapirism. The sequence is believed to grade upwards into the Fountain Spring Beds but the base of the unit is not defined.

Laminated grey pyritic dolomitic siltstones comprising interlayered light grey quartz-dolomite and dark grey biotite-muscovite laminations characterize the unit. The laminations have sharply defined bases with graded tops and commonly display small scale slumping. Quartzites and flaggy, buff weathering,

silty pyritic dolomites occur towards the base of the section. Buff weathering, grey pyritic dolomites, grey quartzites, green shales (ripples and salt casts) and fine sandstones (ripple marks, crosslamination, salt casts) increase in frequency towards the top of the unit.

The laminated light grey-dark grey dolomitic siltstones and silty buff weathering dolomites are also typical of overlying sediments in the lower Fountain Spring Beds, suggesting a gradational boundary but this is uncertain because contacts are faulted.

This unit contains several lithologies typical of those occurring in some of the "Sequences of Uncertain Age", eg. fine sandstones, pyritic dolomites and green shales all with ripple marks and salt casts.

(b) Fountain Spring Beds Eb1

Siltstones, shales and quartzites outcropping on the north-western and eastern margins of the Margaret Inlier are herein defined as the Fountain Spring Beds (Appendix A, Unit 12). Approximately 1 100 m of sediments occur in the type section 3 km north of Mount Margaret. The sequence grades upwards into the Mount Margaret Quartzite, the transition being marked by an increase in orthoquartzites. The base of the Fountain Spring Beds is not exposed but in several localities the unit is observed in tectonic contact with the unnamed siltstone (Eb1). Both sequences are correlated with the River Wakefield Subgroup (Table 1) (Forbes, 1964).

Predominant lithologies include laminated dolomitic meta-siltstones, pale grey quartzite with clay galls, rippled fine sandstones, green slaty siltstones containing salt casts, and minor flaggy buff-weathering grey dolomites near the base.

Quartzite increases at the expense of silt-carbonate upwards through the sequence. A description of the type section is in Appendix D Section 8.

Porphyritic monzonite and syenite sills commonly intrude the Fountain Spring Beds and lower Mount Margaret Quartzite, particularly the siltstone and shale units. Apparently the intrusives were largely unable to penetrate the thick massive quartzites of the upper Mount Margaret Quartzite.

The presence in the Fountain Spring Beds of dolomitic sediments, ripple marks and salt casts suggest marine influence although the upper part of the sequence may have been deposited in a fluvial environment.

(c) Mount Margaret Quartzite Ebo

Reyner (1955) in the text of his report applied the name Mount Margaret Quartzites to a sequence of quartzites he considered to conformably overlie the Duff Creek Beds but he used the name Mount Margaret Formation for the same unit on the published 1-mile geological sheets: Formation was also used by Thomson and Coats (1964). It was Thomson (1969, p. 59) who first used the term Mount Margaret Quartzite. It is proposed in this report that the name Mount Margaret Quartzite apply to a thick sequence of mainly arenaceous sediments conformably overlying the Fountain Spring Beds (Appendix A, Unit 13). The type section, about 5.5 km north-northwest of Mount Fox, includes the complete sequence. The lower contact with the Fountain Spring Beds is gradational and occurs in a section 3 km north of Mount Margaret. The upper contact of the Mount Margaret Quartzite with the Skillogalee Dolomite is conformable. Near the type section there is local brecciation along this upper contact with minor intrusions of carbonate breccia. The large difference in competence between the

thick quartzites of the Mount Margaret Quartzite and the thinly bedded shales and dolomites of the Skillogalee Dolomite probably resulted in this minor brecciation during folding.

The formation is restricted to the southern part of the the Margaret Inlier and it occurs at Mount Margaret itself and also constitutes the Mount Margaret Plateau (Figs 17 & 18). The resistant quartzites forming this imposing landform rise up to 350 m above the surrounding plains.

In the type section, the Mount Margaret Quartzite has a thickness of approximately 2 500 m. It is described in Appendix D, section 9. The sediments are 80% arenaceous and 20% silty-argillaceous. The lower part of the unit consists of interbedded fine slaty quartzitic sandstones (sometimes feldspathic) and green-grey laminated silty shales alternating with massive light grey orthoquartzites. Argillaceous sandstones and thinly bedded cream-grey orthoquartzites with minor interbedded sandy shales (occasionally dark grey) comprise the remainder of the formation. Massive orthoquartzites increase in frequency towards the top of the sequence.

Clay galls, cross-bedding and interference and current ripple marks are common. Mud cracks are rare. It is possible this sequence of mainly arenaceous sediments was deposited in a deltaic environment. Some marine influence, indicated by minor dolomitic siltstones, occurs near the base.

(d) Skillogalee Dolomite Ebk

Conformably overlying the Mount Margaret Quartzite is a sequence containing 3 600 m of dolomites, siltstones, quartzites, sandstones together with minor cherts and magnesites. This sequence



in the Peake and Denison Ranges is far more varied lithologically than the Skillogalee Dolomite in the Clare-Riverton region with which it is correlated (Mirams and Forbes in Thomson et al., 1964).

The formation is confined to the eastern and central parts of the Margaret Inlier and outcrops in a series of basinal synclines (and a complimentary anticline to the southwest) trending north-northwest through the ranges. The formation erodes to a moderately rugged topography but is more subdued than the adjacent quartzites comprising the Mount Margaret Plateau. A preponderance of small trees and shrubs growing on the dolomitic members is a useful criterion in identifying carbonate rich parts of the sequence (Fig. 19). The unit consists of three unnamed members. The basal member conformably overlies the Mount Margaret Quartzite in the vicinity of Mount Fox. The contact is marked by a sudden decrease in elevation from the orthoquartzites to the sandstones and siltstones of the lower Skillogalee Dolomite. This lower sandy member comprises 75% arenaceous sediments, 20-25% argillaceous-silty, 5% dolomitic and has a thickness of 100 m to 3 km northwest of Douglas Well, although the lower contact with the Mount Margaret Quartzite is not preserved here. Main lithologies are platy sometimes gritty sandstones, quartzitic sandstones and green-grey silty shales with rare pale brown, sometimes sandy, dolomites. These mainly arenaceous sediments are characteristic of winnowed sands deposited in beach, shoal and tidal environments. No stromatolites are observed in this member which indicates that the environment was not hospitable to algal-colonization because of the shifting substrate. Precipitation of carbonate was limited to a few thin, pale

dolomites near the top of the member. Clay galls, cross-bedding, graded bedding and ripple marks are common throughout.

This arenaceous member grades into a sequence of dolomites and sandy sediments, the transition being marked by an increased carbonate content (sediments average 55% arenaceous, 15% argillaceous-silty, 30% dolomitic). This middle member reaches a maximum thickness of 1 500 m 17 km north-northwest of Douglas Well. It should be noted that folding has distorted true thickness: both thickening and thinning of the Skillogalee Dolomite are common, especially on the hinges of folds. Most of the dolomites are conglomeratic, cross-bedded and contain reworked dolomite fragments and quartz grit. Colours vary from pale grey-brown to dark grey-black, the latter variety becoming more frequent towards the top of the member. A few stromatolitic dolomite horizons, containing characteristic stromatolites of the group Baicalia, occur in the middle and upper parts of the member. Silicification of the dolomites is common. Associated sediments include green-grey and black pyritic shale, gritty friable sandstone (sometimes feldspathic), thin grey vitreous quartzite, minor blue-black chert and magnesite pebble conglomerate. This member probably records deposition in open lagoons and bays which received considerable amounts of terrigenous sediment. Terrigenous clastic-carbonate couplets imply cyclic sedimentation of unknown complexity but it is probable that upward shoaling, facilitated by seaward outbuilding of carbonate facies, has been followed by rapid transgressions. The vast amounts of reworked dolomite in many of the sediments supports this view. The regressive-rapid transgressive cyclic effects envisaged (due largely to sea level fluctuations) together

with the relatively high clastic content of the sediments explains the paucity of stromatolites. Minor magnesite may have been precipitated by alkaline waters of continental origin reacting with seawater (Forbes, 1961). The magnesite has been reworked into thin, lenticular conglomerates usually less than 0.5 m thick.

The carbonate content shows a marked increase in the upper member and the terrigenous clastic influence is minimal (sediments average 5% arenaceous, 10% argillaceous-silt, 85% dolomitic). Dark grey-black, laminated-thinly bedded dolomites predominate. Reworking of the dolomite into conglomerates is evident and stromatolitic horizons are common. Minor black shale, grey quartzite, feldspathic sandstone, magnesite conglomerate and blue-black chert are interbedded with the dolomites. Deposition probably occurred in shallow cut-off ponds and lagoons with restricted circulation where reducing conditions favoured the accumulation of organic material. Contributions of terrigenous sediments were minimal. The upper boundary of this member is unknown and the estimated maximum thickness is about 1 000 m.

The accumulation of such a large thickness of shallow water sediments (3 600 m) attests to steady basinal subsidence during deposition. In addition, the rapid sedimentation of shallow water carbonates under optimum conditions has resulted in the prevalence of regressive sedimentary cycles. The dolomite component of the Skillsgalee Dolomite accumulated by either direct precipitation of dolomite or formation of syngenetic dolomite by early replacement of calcium carbonate. The second alternative is described by Badiozamani (1973) in a model which requires continuous mixing of groundwaters (low in Mg) with seawater (supplying Mg) during fluctuations of sea level to form a fluid saturated

with respect to dolomite and undersaturated with respect to calcite. During emergence the interface where this mixing occurs becomes a dolomitizing zone and fluctuating sea levels cause this zone to migrate through the sequence.

Preiss, 1973, proposed various environments of deposition for the Skillogalee Dolomite in different parts of the Adelaide Geosyncline. At Adelaide and near Spalding (Bundaleer) the section is 200 m thick and contains between 10 and 30% sandstone which was considered to be deposited in a lagoonal or barred embayment environment. A higher energy environment of deposition occurred in the Willouran Ranges where sandstone forms more than 30% of the section and the section is 1 000 - 4 000 m thick. The northerly trend of increasing sandstone content and increasing thickness are continued in the Peake and Denison Ranges where the section has a minimum thickness of 3 600 m and contains 40-50% sandstone, mostly concentrated in the lower two members. The increased thickness and terrigenous clastic component reflects active tectonism in the northern portion of the Adelaide Geosyncline with continental sources existing to the west and southwest.

(e) Unnamed transition unit Eb2

In an anticlinal structure on the southwest margin of Margaret Inlier, 6 km north of Box Creek R.S., 3 000 m of upper Burra Group sediments disconformably underlie the Sturtian tillite in a section referred to as the Boorthanna Section (Fig. 20). Sediments are predominantly thinly bedded, fine-grained clastics with minor, locally abundant, carbonates (sediments average 60% arenaceous, 25% argillaceous-silty, 15% dolomitic). The upper part of the section was originally described by Cooper et al.,

(1964). Interest was rekindled in the Boorthanna Section following discovery of a well-preserved late Precambrian microflora in stromatolitic black cherts collected by H. Wopfner (Schopf and Fairchild, 1973). Later Fairchild (1975) subdivided the sequence into seven units (units A-G, Fig. 20). This is a valid lithological subdivision which forms the basis of the following description. Units A-D comprise the lower 2 000 m of the Boorthanna section and represent the unnamed unit which is transitional between the underlying Skillogalee Dolomite and overlying Kalachalpa Formation (Units E-G of Fairchild, (1975)).

Unit A consists of 1 500 m of mainly laminated, gritty, quartzitic sandstones (often feldspathic), interbedded grey-green shales with minor laminated grey-brown dolomites (sediments average 75-80% arenaceous, 20% argillaceous-silty, 0-5% dolomitic).

Unit B is more dolomitic (sediments average 25% arenaceous, 35% argillaceous-silty, 40% dolomitic) and consists of 190 m of alternating dolomites and fine clastics with black chert interbeds and minor magnesite pebble conglomerates. Dolomites are conglomeratic and often silicified and several stromatolitic horizons are present (including Baicalia).

The clastic content increases in Unit C (sediments average 90% arenaceous, 10% dolomitic) which is predominantly platy friable sandstones with minor magnesite pebble conglomerates, orange dolomites and silicified carbonate pebble conglomerates.

Unit D contains characteristic coarse chert pebble conglomerates

(Fig. 21) magnesite pebble conglomerates, black cherts and stromatolitic dolomites (sediments average 45% arenaceous, 15% argillaceous-silty, 40% dolomitic).

Mud cracks, ripple marks, cross-bedding, graded bedding and convolute bedding are common throughout and suggest a shallow marine environment of deposition which was intermittently exposed and desiccated. There is extensive evidence of penecontemporaneous reworking (e.g. intraformational conglomerates and erosional micro-unconformities (Fig. 22). Cyclic sedimentation appears to be similar to that operating during the deposition of the Skillogalee Dolomite and the mode of dolomitization is also probably similar. Fairchild (1975) suggests silicification may be related to regressive cycles during which evaporative pumping of silica-enriched groundwaters facilitated silicification.

Units A-D have a number of affinities with the Skillogalee Dolomite which include the presence of;

- 1) magnesite pebble conglomerates,
- 2) stromatolites of the group Baicalia. Fairchild (1975) discovered stromatolitic microfloras in eighteen beds in the Boorthanna section,
- 3) chert, conglomeratic dolomite together with other lithologies and sedimentary structures indicating a similar environment of deposition to that of the Skillogalee Dolomite.

However, the unnamed sequence is not as dolomite-rich as the Skillogalee Dolomite (Skillogalee Dolomite sediments average 35-40% dolomitic, 45-50% arenaceous, 15% argillaceous-silty; unnamed unit averages 10-15% dolomitic, 70% arenaceous, 15-20% argillaceous-silty). The Skillogalee Dolomite is less extensively silicified and contains only minor chert. No stratigraphic contact between the two units has been observed and there is no definite lithological

overlap recognised between the upper Skillogalee Dolomite and the lower part of the unnamed unit. Thus, barring rapid facies changes, it is apparent that part of the upper Torrensian sedimentary record is missing, the magnitude of the gap being uncertain. The fact that the unnamed unit has many lithological affinities with the overlying Kalachalpa Formation as well as the underlying Skillogalee Dolomite indicates that it is transitional between these two formations.

An equivalent sequence, faulted against the Kalachalpa Formation, outcrops 1 km east of the Box Creek R.S. It is doubtful if the lithological subdivision of Fairchild (1975), i.e. Units A-D, applies here as the upper part of the sequence is considerably less sandy than its counterpart to the north.

(f) Kalachalpa Formation Ebh

The upper 900 m of sediments outcropping in the Boorthanna Section, 6 km north of Box Creek R.S. are herein named the Kalachalpa Formation (Appendix A, Unit 11) which is composed mainly of fine grained clastics and associated dolomite horizons (Appendix D, Section 10). The sequence corresponds to units E-G of Fairchild (1975) (Fig. 20) and conformably overlies the unnamed transitional unit. The upper contact with the Sturtian glacials is marked by a disconformity. The Kalachalpa Formation correlates with the Myrtle Springs Formation (Coats, 1973) (Table 1).

Although the Kalachalpa Formation has lithological affinities with the conformably underlying unnamed transitional unit (units A-D of Fairchild) it differs in a number of respects. The Kalachalpa Formation;

- 1) is more argillaceous, containing 45-50% arenaceous, 35% argillaceous and 15-20% dolomitic sediments,
- 2) contains a higher proportion of oolitic sediments,
- 3) contains only a minor percentage of chert,

- 4) does not contain any magnesite conglomerates,
- 5) records a wide variety of stromatolites (Fig. 23; Fairchild, 1975).
- 6) does not contain silicified dolomite pebble conglomerates or chert pebble conglomerates. Fairchild (1975) suggests silicification may have occurred late in the diagenetic history.

The most useful criteria for identifying the contact between the unnamed transitional unit and the Kalachalpa Formation in the field are;

- a) the last occurrence of chert pebble conglomerates,
- b) an increase in frequency of green-grey, olive green and occasionally dark red-brown argillaceous units,
- c) a decrease in the amount of dolomite from unit D to unit E.

The basal part of the Kalachalpa Formation, Unit E, consists of 365 m of platy siltstones and fine sandstones with minor dolomites (sediments average 55% arenaceous, 45% argillaceous, 5% dolomitic). There is one distinctive biohermal stromatolite horizon. Dolomite content increases in unit F (sediments average 35% arenaceous, 30% argillaceous, 35% dolomitic). Slaty red and green siltstones and a few orthoquartzites separated by platy sandstones predominate. The dolomites are often sandy and become increasingly stromatolitic towards the top of the unit. Black chert and oolitic sediments are locally abundant. Total thickness is 380 m. Unit G is made up of 175 m of mainly arenaceous sediments (sediments average 75% arenaceous, 25% argillaceous). Main lithologies are orthoquartzites and sandstones with minor slaty and sandy siltstones.

Mud cracks, clay galls, ripple marks, cross-bedding, intra-formational conglomerates etc. indicate deposition in a shallow water environment similar to that existing during deposition of units A-D.



A study of palaeocurrent indicators by Fairchild (1975) indicates sediment transport was generally from west to east and southeast with the Gawler Craton constituting the principal source area. The predominance of clastic sediments in the Boorthanna section is consistent with proximity of the Boorthanna area (and the Peake and Denison Ranges generally) to active continental sources probably situated to the west and southwest.

#### UMBERATANA GROUP (STURTIAN)

In the Peake and Denison Ranges there was a period of mild erosion prior to the onset of Umberatana Group sediments, with lower Sturtian glacials deposited disconformably over upper Burra Group sediments on the northern limb of the Box Creek Syncline. There is no evidence of significant Sturtian tectonism. The following units comprise the Umberatana Group in the Peake and Denison Ranges - Calthorinna Tillite (new name), unnamed sandstone, Tapley Hill Formation, Thora Dolomite (new name) and an unnamed siltstone belonging to the Willochra Subgroup (Table 10).

TABLE 10

#### Stratigraphy - UMBERATANA GROUP

Ph1	Unnamed siltstone	} Willochra Subgroup
Bf1	Thora Dolomite	
Bft	Tapley Hill Formation	} Farina Subgroup
Bul	Unnamed sandstone	
Pub	Calthorinna Tillite	

#### BURRA GROUP - Kalachalpa F.

————— conformable contact

——— interfingering contact

——— disconformity

The Umberatana Group as defined by Coats (1964) consists of a lower (Sturtian) and upper (Marinoan) glacial phase separated by an interglacial period. However, the lower glacial phase is now considered to consist of two glacial episodes separated by an unconformity (Coats, 1973). The Sturtian glacials (Calthorinna Tillite and unnamed sandstone) are thought to be representative of the second episode of the lower glacial phase on the basis of gritty quartzite clasts, with fossil red weathering, assumed to have been reworked from the Pandurra Formation. This criterion has been used elsewhere in the Adelaide Geosyncline to distinguish between the first and second episodes of the lower (Sturtian) glaciation (Coats and Forbes, 1977). An unnamed sandstone unit overlies and interfingers with the Calthorinna Tillite and marks the end of the glacial period. There followed a marine transgression with the deposition of marine sequences comprising the Tapley Hill Formation, Thora Dolomite and an unnamed siltstone.

Umberatana Group sediments are restricted in outcrop to a syncline 1 km northeast of Box Creek R.S. (Box Creek Syncline) and a rafted block in a diapir 6 km south of Mount Anna. Approximately 2 200 m of Sturtian sediments outcrop on the southern limb of the Box Creek Syncline.

(a) Calthorinna Tillite Eub

Disconformably overlying the Kalachalpa Formation is a sequence of glaciomarine and marine sediments herein named the Calthorinna Tillite (Appendix A, Unit 15). The type section, 5.5 km northwest of Box Creek R.S., consists of approximately 650 m of diamictites, conglomerates and interbedded marine sediments (Appendix E, Section 11).

Dominant lithologies comprising the Calthorinna Tillite are gritty quartzitic sandstones, medium-coarse feldspathic sandstones, thinly bedded pale yellow-grey pebbly dolomites and dolomitic siltstones, finely laminated pale grey-green shales plus interbedded diamictite and conglomeratic dolomites (Figs 24, 25 & 26). The formation is characterized by rapid lateral and vertical facies changes. The conglomerate and diamictite units are variable in composition, texture and thickness. The conglomerates are massive, buff coloured dolomites containing pebbles and cobbles of mainly chert and dolomite and minor quartz grit. A few exotic clasts occur in these conglomerates which are usually less than 10 m thick.

The diamictite units are variable in thickness up to 100 m. The matrix contains varying proportions of quartz, dolomite and clay with the silt-size fraction predominating. Bedding is occasionally observed. Pebble and granule size clasts predominate with cobbles less common and boulders rare.

Clast lithologies have a dominant local component but there is considerable evidence for more distal southwesterly sources including the Mount Woods Inlier on BILLA KALINA and the Gawler Ranges. Main clast lithologies and suspected sources are listed below;

- a) oolitic chert, pale and dark grey laminated chert, silicified chert and quartz pebble-conglomerate, dark grey and pale yellow dolomite, siltstone and stromatolitic dolomite (Burra Group sediments - local source),
- b) red porphyritic granite and hornblende granite (Balta Granite-Mount Woods Inlier),

- c) red porphyritic rhyolite (Gawler Range Volcanics),
- d) rare clasts of vesicular basalt (Cadlawarena Volcanics-local source),
- e) numerous types of quartzite and quartzitic sandstone with one boulder of gritty quartzite showing fossil red weathering (?Pandurra Formation - R.P. Coats, pers. comm),
- f) dolomitic sandstone and laminated pale yellow dolomite (Duff Creek Beds - local source),
- g) microgranodiorite, dark grey schist and phyllite (source unknown).

In addition to the sources listed above it is probable a small percentage of dolomite and chert clasts in the diamictite are intraformational. Conglomeratic dolomites contain a high proportion of intraformational clasts (etched pale green and grey dolomite and cream to dark grey chert) presumably reworked during rapid transgressive - regressive cycles as sea level responded to temperature fluctuations (Fig. 24).

On the northern limb of the Box Creek Syncline a basal green-brown diamictite, about 100 m thick, disconformably overlies the upper part of the Kalachalpa Formation which consists of mainly quartzites with clay gall laminations.

The matrix of the diamictite is usually silty although it becomes more quartzitic towards the base of the unit. A silicified channelfill conglomerate containing mainly chert pebbles and a cross-bedded sandstone occur near the base in a railway cutting 4.5 km northwest of Anna Creek R.S. A fissile, olive green shale interfingers with the diamictite.

Overlying this basal unit are 500 m of gritty and pebbly sandstones, conglomeratic dolomites, dolomitic silty shales, quartzites and chocolate-green shales with diamictite interbeds. The uppermost unit observed in this northern section is a pink-brown, massive conglomeratic dolomite containing mainly chert and pale dolomite fragments and clasts of green matrix diamictite presumably reworked from lower in the section (Fig. 26). In the upper part of the section gritty feldspathic sandstones, typical of the overlying unnamed sandstone (Pul) interfinger with diamictites and conglomerates of the Calthorinna Tillite.

On the southern limb of the Box Creek Syncline the Calthorinna Tillite exhibits faulted contacts with Burra Group sediments. A comprehensive section is located 1.5 km northeast of Box Creek R.S. where 450 m of diamictites, conglomerates and interbedded marine sediments outcrop (Appendix E, Section 12). The base of this section is marked by a cross-bedded quartzite containing flat shale pebble conglomerate laminations. This unit is sheared and silicified indicating a tectonic contact with underlying stromatolitic dolomites of the Kalachalpa Formation. An unnamed sandstone normally overlies the Calthorinna Tillite on this southern limb of the syncline. There is no apparent interfingering between these two units as observed on the northern limb of the Box Creek Syncline and in a rafted block in a diapir 6 km south of Mount Anna.

Sediments of the Calthorinna Tillite were deposited under glacial marine and marine conditions. It is unlikely the diamictites were deposited from a grounded continental ice sheet considering their lenticular nature and in many cases carbonate-rich matrix.

The presence of intimately associated marine sediments supports this view. Quartz and clay-rich diamictites may have been deposited from a wet-based glacier grounded offshore (Carey and Ahmad, 1960). Other diamictite interbeds, including the more carbonate-rich varieties, may represent reworked till deposited by mudflows and debris flows (turbidity currents) on a foreset slope where the glacier became buoyant. There may be minor contributions of ice-rafted material.

Interbedded with mixtites are dolomites, shales and arenaceous sediments displaying ripple marks, cross-laminations, and graded bedding which were deposited under marine conditions as sea level responded to temperature fluctuations. Some of the laminated shales resemble varves, and rare clusters of boulders and cobbles suggest ice rafting during periods of high sea level. Conglomeratic dolomites (containing almost solely chert and dolomite fragments and in one case clasts of silty green-matrix diamictite) represent intra-formational conglomerates formed by cannibalization of earlier sediments during periodic sea level fluctuations (Fig. 24). Evidence of cyclic sedimentation is most convincing on the southern limb of the Box Creek Syncline where conglomeratic dolomite-mixtite-sandstone triplets are repeated through the sequence.

As discussed by other workers (eg. Spencer, 1971; Young, 1976) the intimate association of glacial rocks with dolomitic sediments, which normally are considered indicative of warm climate and low latitude, is incompatible. This anomalous association occurs in many parts of the world and Young (1976) summarises explanations provided by various writers. Two of the more attractive theories are:

- a) Williams (1975) suggested that secular variation in obliquity of the earth's ecliptic gives rise to periodic low latitude glaciation and extreme seasonal variation in climate;
- b) differences in the earth's atmosphere in the Precambrian (high  $\text{PCO}_2$ ), as suggested by Young (1976), permitted sedimentation of rock types normally ascribed to a warm climate regime in relatively high latitudes subject to periodic glaciation.

(b) Unnamed sandstone Bul

Conformably overlying and interfingering with the Calthorinna Tillite is a sequence of feldspathic sandstone and quartzitic sandstone. The type section, 2 km east-northeast of Box Creek R.S., consists of 530 m of medium to coarse-grained, well-rounded, feldspathic sandstone displaying large scale cross-bedding up to 4 m in height. Subordinate lithologies include thickly bedded quartzitic sandstones (sometimes feldspathic) with minor micro-conglomerates and argillaceous sandstones. A detailed description of this section is an Appendix E, Section 12.

On the southern limb of the Box Creek Syncline this sandstone unit maintains a constant thickness along strike for about 6 km. On the northern limb a number of gritty feldspathic sandstones are interbedded with diamictites, conglomerates and shales illustrating interfingering between the sandstone unit and the Calthorinna Tillite.

Deposition of this unit marks the end of the glacial period. As temperatures rose so did sea levels, and meltwater streams probably transported a large amount of sorted glacial material into a

deltaic-marine environment where the thick sandstone unit was deposited. Large scale cross-beds indicate a high energy environment of deposition and the abundance of fresh feldspar suggests that cold conditions still prevailed. Interfingering of the sandstone and Calthorinna Tillite on the northern limb of the Box Creek Syncline supports this view and indicates flagging glacial activity prior to the major marine transgression associated with the deposition of the Tapley Hill Formation. Approximately 2.5 km northeast of Box Creek R.S. some of the sandstone beds appear to be truncated at a shallow angle by shales of the overlying Tapley Hill Formation. These mesoscopic trends may represent large foreset beds of glacial outwash material, deposited in a deltaic-marine environment, which have been truncated by the overlying transgressive marine sequences. Red granules of porphyry are common throughout the sandstone. They may have been reworked locally from clasts in the tillite although a more southwesterly origin from the Gawler Range Volcanics is a possibility.

(c) Tapley Hill Formation Eft

The Tapley Hill Formation, which comprises a consistent facies throughout the Adelaide Geosyncline (Coats, 1971), is slightly more calcareous and much reduced in thickness in the Peake and Denison Ranges compared with the type sequence in the Mount Painter Province (Coats, in Thomson et al., 1964). The formation outcrops only on the southern limb of the Box Creek Syncline where a total thickness of 150 m is recorded. A description of this section is in Appendix E, Section 12.

The deposition of this formation marks the onset of marine conditions following termination of glacial activity. The base of this transgressive sequence occasionally truncates bedding planes in the underlying sandstone unit. At the base is a 10 m-thick pale yellow dolomite, containing rounded quartz grains, which thins



to the east. The overlying 140 m consists of finely laminated dark grey to black silty shales and silty dolomites which weather grey-green. The sequence becomes more dolomitic towards the top.

(d) THORA DOLOMITE Bf1

Conformably overlying the Tapley Hill Formation is a distinctive dolomite, 30-40 m in thickness, herein named the Thora Dolomite (Appendix A, Unit 16). The type section is on the southern limb of the Box Creek Syncline and a description is in Appendix E, Section 12.

The unit consists of red-brown to orange weathering grey-green to brown dolomite with minor interbedded siltstone. Bedding is irregular and ranges up to 20 cm in thickness. Algal bedding and large festoon surfaces are common. A few intraformational dolomite fragments suggest minor reworking of the dolomite in a shallow marine environment. The Thora Dolomite is correlated with the Brighton Limestone of Mawson and Sprigg (1950).

(e) Unnamed Siltstone Bh1 (Marinoan)

Conformably overlying the Thora Dolomite are 950 m of mainly green-grey and reddish silty shales and fine sandstones. The thickest section is on the southern limb of the Box Creek Syncline, 5 km east-northeast of Box Creek R.S. (Appendix E, Section 12).

The Willochra Subgroup as defined by Thomson (in Parkin, 1969) is characterised by reddish, medium - coarse grained clastics with minor tongues of granule limestone. This unnamed siltstone records the first appearance of red beds in the Umberatana Group in the Peake and Denison Ranges and hence the unit is assigned to the Willochra Subgroup.

The base of the unit consists of grey-green siltstones, dolomitic siltstones and fine argillaceous sandstones. The major part of the sequence consists of interbedded finely laminated grey-green and red-brown silty shales with ripple marks, fine sandstones, thin

(50 cm) grey-yellow dolomites and silty dolomites. The dolomites are frequently silicified and occasionally gritty and the sand content generally increases in the upper part of the sequence with several thin reddish coarse sandstones occurring towards the top. The red-brown colour and gradual coarsening upwards of the sediments indicates a regressive sequence. The upper boundary of the unit is not exposed.

#### BUNGADILLINA MONZONITE      Odb (ORDOVICIAN)

An igneous suite of monzonites, syenites, diorites, albitites and minor granites outcropping 12 km northeast of Nilpinna Homestead are collectively defined as the Bungadillina Monzonite (Encl. 1). The cluster of intrusive bodies is coincident with a major lineament (interpreted from E.R.T.S. Landsat imagery) which is an extension of the Karari Fault and which trends SW-NE, intersecting the ranges where the monzonitic suite outcrops (Fig. 2). K-Ar geochronology yields a Delamerian age for intrusion of the Bungadillina Monzonite.

In the two largest intrusive bodies northeast of Nilpinna Homestead, porphyritic monzonites and syenites are the dominant rock types (Fig. 27) while porphyritic diorites are less frequent. Feldspar phenocrysts are ubiquitous and often define a mineral elongation trending  $155$  to  $185^{\circ}$  (Fig. 28). The feldspars are euhedral and many are 30 mm in length. Potash feldspar phenocrysts are dominantly perthitic orthoclase and microcline while plagioclase phenocrysts are strongly zoned with sericitic cores and clear rims (Fig. 29). Water-clear plagioclase rims mantling potash feldspar crystals have also been recorded (Turner, 1969). Plagioclase compositions (Turner, 1969) fall in the range

An<sub>30-40</sub>. Free quartz is often absent, but if present forms less than 10% of the rock. Augite, hornblende and biotite are locally abundant and may rarely constitute 25% of the rock. Accessory

minerals are magnetite, garnet, sphene, epidote and calcite.

In the diapiric zone in the headwaters of Edwards and Bulldog Creeks are numerous, small, irregularly-shaped intrusives. Albitites are common within this zone and consist of greater than 95% plagioclase. In the accompanying diorites, hornblende and plagioclase are dominant constituents with many euhedral hornblende crystals, up to 10 mm across, mantled by plagioclase. Microgranites, granodiorites and leucoadamellites are rare and when present form small circular stocks.

Porphyritic monzonites and syenites intruding the Fountain Spring Beds and Mount Margaret Quartzite have formed a series of sills up to 6 km long and 20 m thick. Numerous xenoliths of quartz-chlorite schists, metasiltstones and rounded quartz grains indicate considerable contamination by nearby Adelaidean sediments. Phenocrysts within the sills are hornblende and zoned potash and plagioclase feldspars, within a fine grained felsic groundmass. Large plagioclase phenocrysts are mantled by potash feldspar (Steveson, 1976 b; 1976 c).

Silicate analyses of two specimens, a hornblende syenite and hornblende monzonite are tabulated in Table 11. The analyses reveal a significant difference from the average for monzonites and syenites quoted by Le Maitre (1976). These include an enrichment in  $\text{SiO}_2$  at the expense of  $\text{TiO}_2$ ,  $\text{FeO}$ ,  $\text{MnO}$ ,  $\text{CaO}$  and  $\text{P}_2\text{O}_5$ .

TABLE 11  
BUNGADILLINA MONZONITE

	1	2	3	4
SiO <sub>2</sub>	65.25	58.58	65.36	62.60
TiO <sub>2</sub>	.24	.84	.28	.78
Al <sub>2</sub> O <sub>3</sub>	16.26	16.64	17.41	15.65
Fe <sub>2</sub> O <sub>3</sub>	1.30	3.04	1.52	1.92
FeO	1.00	3.13	1.20	3.08
MnO	.02	.13	.03	.10
MgO	.59	1.87	.60	2.02
CaO	1.85	3.53	2.66	4.17
Na <sub>2</sub> O	4.65	5.24	5.22	3.73
K <sub>2</sub> O	6.76	4.95	4.55	4.06
P <sub>2</sub> O <sub>5</sub>	.09	.29	.09	.25
H <sub>2</sub> O <sup>+</sup>	.24	.99	.39	.90
H <sub>2</sub> O <sup>-</sup>	.19	.23	.17	.19
TOTAL	98.44	99.46	99.47	99.45

1. Hornblende syenite (Bungadillina Monzonite) northeast of Nilpinna Homestead. Specimen 6141 RS 18.
2. Average of 102 syenites (Le Maitre, 1976)
3. Hornblende monzonite (Bungadillina Monzonite) north-northeast of Nilpinna Homestead. Specimen 6041 RS 58.
4. Average of 102 monzonites (Le Maitre, 1976).

Seven monzonites and syenites and one dolerite dated by the K-Ar method on hornblende, biotite, potash feldspar and total rock yielded ages ranging from 459 to 667 m.y. (Webb, 1970; Webb & Lowder, 1971; 1972). The Bungadillina Monzonite is interpreted as being a late-stage orogenic intrusive of the Delamerian Orogeny

(circa 500 m.y.) and is analogous to similar granitic intrusives over a wide area in the southeastern part of the State (Webb, 1976).

Fission track dating on apatite on one specimen (horn-blended syenite - 6141 RS 18) yielded a Permian age of  $266 \pm 23$  Ma (average of six apatite grains) (Radke, 1973) and may be interpreted as indicating tectonism during the Permian.

### Dykes

In the southern part of the ranges several dolerite and uralitized dolerite dykes intrude dolomites, sandstones and siltstones of the Skillogalee Dolomite and Duff Creek Beds (Encl. 1). The dykes have long, linear outcrops which suggest a late-stage orogenic time of intrusion. The dolerite dykes have undergone varying degrees of alteration. Turner (1969) reported olivine, augite and plagioclase in a dolerite to the northeast of Nilpinna H.S. while Cooper (1976) recorded abundant quartz, chlorite, dolomite and muscovite in altered dolerites near Davenport Creek.

Near a crocidolite-malachite mine 2 km west of Tarlton Springs are rare and unusual spherulitic dykes up to 0.3 m wide. The spherulites which are up to 5 mm in diameter are composed of radiating potassium feldspar crystals in a secondary dolomite and quartz matrix (Steveson, 1976b). Spherulites have been interpreted to be the product of devitrification of glassy lavas with the spherulites being enriched in potassium as the degree of spherulite development increases (Ewart, 1971). The spherulitic potassium feldspar dyke near Tarlton Springs is believed to be a late stage acidic intrusive of the Bungadillina Monzonite.

## BOORTHANNA FORMATION P1b (PERMIAN)

Scattered outcrops of Permian sediments along the western margins of the Peake and Denison Ranges and Dutton Inlier were first recognised by Reyner (1955). Townsend and Ludbrook (1975) named these sediments the Boorthanna Formation and considered the abundance of striated, faceted and soled pebbles to indicate a glacial origin.

A number of small outcrops of Permian sediments occur several kilometres northeast of Box Creek R.S. Some of the outcrops consist of shallow dipping, gritty dolomites with occasional boulders. Several areas of closely scattered boulders (many of which are striated and faceted) represent lag deposits (Fig. 30). Similar rocks are observed 5 km northwest of Duff Creek R.S. A small pocket of consolidated Permian diamictite outcrops 6.5 km north of Warrina R.S. Striated and faceted clasts indicate a glacial origin: local Adelaidean rocks comprise the dominant clast lithologies and clasts derived from Burra Group and Umberatana Group sediments predominate.

Permian sediments outcropping on the western and southwestern margins of the Dutton Inlier are described in detail by Heath (1965). According to Heath the Permian section is at least 35 m thick, with the base transgressive onto Adelaidean rocks. The lower part of the sequence consists of gritty dolomites (1 m thick) which grade into diamictites (30m thick) and bluish laminated shales. The Permian sediments outcrop poorly; erosion has resulted in extensive coarse lag deposits of faceted and striated boulders of a wide variety of rock types. Some of the pebbles and boulders, particularly the red porphyries, may have originated by erosion of the overlying conglomeratic and boulder beds of the Algebuckina Sandstone and Cadna-owie Formation (Heath, 1965).

Ludbrook (1961) presented lithological and palaeontological data from available cores and sludges from Boorthanna and Anna Creek Bores drilled in 1911 and 1887 respectively. Boorthanna Bore, 14 km southeast of Nilpinna H.S., penetrated nearly 590 m of Permian sediments without disclosing the base. Anna Creek Bore, 18.5 km south of Nilpinna H.S. records approximately 300 m of Permian sediments lying on Precambrian basement. Grey carbonaceous siltstones, mudstones and occasional sandstones are the most prominent lithologies. Smith (1973), following a study of gravity data on the southwestern flank of Margaret Inlier, discussed a prominent negative gravity anomaly passing through the vicinity of Boorthanna and Anna Creek Bores. Smith notes this negative gravity anomaly circles the southern margin of the ranges passing between Jurassic and Precambrian outcrops and suggests the feature may record a locally preserved Permian sequence at least 60 m thick.

DIAPIRISM

Diapirism is common throughout the Peake and Denison Ranges and plays an important role in the present geological framework of the region. Diapiric activity is recorded in two general contexts;

- a) broad zones of disruption containing disorientated, rafted blocks of varying size and lithology incorporated in a carbonate-matrix breccia,
- b) narrow zones of carbonate breccia intruded as sills, plugs and more rarely as dykes. Diapiric breccia is often intruded along faults and anticlinal fold hinges.

A broad diapiric zone, several kilometres wide, occurs above the Cadlareena Volcanics on Warrina (Encl. 1). Incorporated within the diapiric breccia are rafted blocks of quartzites, siltstones, dolomites, cherts and occasional volcanics mostly derived from

lower Adelaidean sequences. Several blocks contain quartzites of the Baltucoodna Quartzite and one contains basement quartzites unconformably overlain by Younghusband Conglomerate and Cadlarena Volcanics.

The blocks are up to several kilometres across and bedding is often highly contorted, and incompetent shale and dolomite units display extremely plastic deformation. Descriptions of some of the larger blocks are on pages 42 & 43. The carbonate-matrix breccia, in which these blocks are incorporated, weathers easily and forms a subdued topography. The diapiric zones are often characterized by a conglomeration of resistant, disorientated blocks, possessing characteristic contorted bedding, set in lower lying and poorly outcropping carbonate breccia (Figs. 31 & 32).

In the diapiric zone above the Cadlarena Volcanics on Warrina, the degree of disruption decreases away from the volcanics to the southwest.

A transition is seen from:-

- a) a zone of complete disruption; diapiric breccia is common and blocks of contrasting lithologies are observed in confused orientations;
- to b) a zone of less intense disruption; minor diapiric breccia and dislocation of thick sequences (Murrana Beds) but with retention of original bedding orientations;
- to c) a generally undisturbed stratigraphic sequence (Murrana Beds) with only minor tongues and plugs of diapiric breccia.

A second large diapiric zone occurs in the headwaters of Levi Creek about 16 km east of Duff Creek R.S. This disrupted zone, 2-3 km wide represents the intersection of two major anticlinal fold hinges. Faulting, intrusion of carbonate breccia, brecciation and plastic deformation are characteristic. Intrusive



lower Adelaidean sequences. Several blocks contain quartzites of the Baltucoodna Quartzite and one contains basement quartzites unconformably overlain by Younghusband Conglomerate and Cadlareena Volcanics.

The blocks are up to several kilometres across and bedding is often highly contorted, and incompetent shale and dolomite units display extremely plastic deformation. Descriptions of some of the larger blocks are on pages 42 & 43. The carbonate-matrix breccia, in which these blocks are incorporated, weathers easily and forms a subdued topography. The diapiric zones are often characterized by a conglomeration of resistant, disorientated blocks, possessing characteristic contorted bedding, set in lower lying and poorly outcropping carbonate breccia (Figs. 31 & 32).

In the diapiric zone above the Cadlareena Volcanics on Warrina, the degree of disruption decreases away from the volcanics to the southwest.

A transition is seen from:

- a) a zone of complete disruption; diapiric breccia is common and blocks of contrasting lithologies are observed in confused orientations;
- to b) a zone of less intense disruption; minor diapiric breccia and dislocation of thick sequences (Murrana Beds) but with retention of original bedding orientations;
- to c) a generally undisturbed stratigraphic sequence (Murrana Beds) with only minor tongues and plugs of diapiric breccia.

A second large diapiric zone occurs in the headwaters of Levi Creek about 16 km east of Duff Creek R.S. This disrupted zone, 2-3 km wide represents the intersection of two major anticlinal fold hinges. Faulting, intrusion of carbonate breccia, brecciation and plastic deformation are characteristic. Intrusive

albitites, diorites, syenites and monzonites are concentrated in this zone and photointerpretation of LANDSAT 1 imagery reveals this zone lies along a lineament which is an extension of the Karrari Fault (Fig. 2). This disrupted anticlinal zone reflects movements triggered deep in the crust during the Delamerian Orogeny.

Disruption always occurs immediately above the Cadlareena Volcanics. As seen elsewhere in the Adelaide Geosyncline, lower Adelaidean sequences have been most severely affected by diapirism (Thomson and Coats, 1964; Coats, 1973).

Diapiric carbonate is intruded along many faults (e.g. along the fault marking the eastern margin of the ranges near Peake ruins). Breccia is also often intruded parallel to bedding in relatively undisturbed sequences as is the case in the lower Murrana Beds 4 km east of Warrina R.S. Tongues, dykes, sills and plugs of diapiric breccia commonly intrude deformed and block-faulted sequences as seen on the eastern side of Margaret Inlier, south of Levi Creek.

Several small pockets of brecciated Adelaidean sediments set in carbonate breccia are entirely surrounded by basement rocks to the south of Mount Kingston and also on the southern margin of the Denison Inlier. These small outliers may represent remnants of Adelaidean rocks thrust over basement rocks. The southern margin of the Denison Inlier is bound by brecciated Adelaidean rocks and carbonate breccia which intrudes basement rocks along faults, indicating high mobility for the diapiric material.

The source of the carbonate breccia is uncertain. In the Mount Painter Province the Wywyana Formation (actinolitic marble unit) can be traced from normal stratigraphic sequence into diapiric context and it has been suggested this formation is the source material for diapirs in the Adelaide Geosyncline (Coats, 1971; 1973). The equivalent of this formation in the Peake and Denison Ranges is the

Coominaree Dolomite, which may be the source of the diapiric breccia but there is no conclusive evidence to support this view. On the contrary, the Coominaree Dolomite has behaved in a competent manner during deformation where it is seen in undisturbed sequence on the margins of the Coominaree Mine Block and War Loan Mine Block. Nowhere was this unit traced from stratigraphic sequence into diapiric context. It is significant that disruption occurs immediately above the Cadlareena Volcanics in a number of localities suggesting the source of diapiric breccia may have been higher in the sequence above the volcanics. It should be noted however, that disruption has occurred below the volcanics; notably 0.5 km northeast of War Loan Mine and on the western margin of the Rockwater Hill Block, where decollement with basement rocks was associated with intrusion of diapiric carbonate breccia.

Salt casts are locally abundant in lower Adelaidean rocks and many blocks in the diapir are enriched in salt casts relative to adjacent sequences. However, there is no evidence of evaporite minerals in the carbonate breccia, and there are no evaporite beds recorded in lower Adelaidean sequences. Consequently the role that salt has played in diapirism is uncertain. It is possible that sequences closely overlying the volcanics and containing evaporites provided a source for the diapirs. All evidence of such source beds would be obliterated by diapiric processes. The apparent absence of evaporite strata led Dalgarno and Johnson (1968) to the assumption "that extreme incompetence of the Callanna Beds under stress caused first the decollement with the basement and then piercement folding."

On COPLEY, evidence from the Burr Diapir suggests that the earliest diapirism occurred prior to deposition of the Umberatana Group (Coats, 1973). Subsequent to a study of the Blinman Diapir

Coats (1965) suggested diapirism may have been initiated by a mild orogeny prior to the Sturtian Glaciation and continued through to Cambrian times. There is no evidence of significant Sturtian tectonism in the Peake and Denison Ranges, and there is no indication of diapiric detritus in Adelaidean sediments which makes dating of diapiric activity difficult. The youngest Adelaidean rocks outcropping (Umberatana Group) are intruded by diapiric breccia and it appears that the major phase of diapirism was synchronous with the Delamerian Orogeny (late Cambrian to early Ordovician).

The main evidence supporting this view is;

- 1) presence of blocks of Ordovician syenite in a diapir,
- 2) presence of strongly folded and cleaved blocks in diapirs,
- 3) the proliferation of carbonate breccia intruded along many faults,
- 4) presence of disrupted anticlinal hinge zones.

It is noteworthy that leucogranite stocks of Ordovician age intrude diapirs in several localities e.g. in the diapiric zone at the head of Levi Creek.

#### TECTONISM

Rocks within the Peake and Denison Ranges display a complex deformational history. At least six periods of tectonism have been recognised ranging in age from Carpentarian to Quaternary.

The first deformation is evidenced by a layer-parallel foliation in units of the Peake Metamorphics. This is expressed by a muscovite, biotite, or chlorite schistosity and gneissosity in the metasediments and a hornblende - chlorite schistosity in metamorphosed diorites and basalts. The Wirriecurrie Granite has a biotite schistosity which is parallel to the layer-parallel schistosity of adjacent metamorphics, implying granite intrusion synchronous with the first deformation episode. Rb-Sr geochronology on the granite and K-Ar dating on diorite

sills (Webb & Lowder, 1972; Webb, 1974) indicate a Carpentarian age (1600 Ma) for the tectonism.

The second tectonic phase is indicated by the updating of geochronological dates on the Wirriecurrie Granite. Both Rb-Sr and K-Ar methods suggest an orogenic event about 1050 Ma, which probably equates with the Musgravian Orogeny. No faults or folds have been recognised as being of this generation.

Tectonism during the Delamerian Orogeny has resulted in folding, faulting and disruption of Adelaidean sediments and refolding of basement metamorphics. The widespread diapirism was probably coincident with this orogenic phase. Cross folding with approximately north-south and east-west striking axial planes have formed open basins and tight, disrupted domes in Adelaidean sediments. These are exemplified by the broad elongate basin of Skillogalee Dolomite south of Mount Margaret and a disrupted, elongate dome just north of Anna Creek railway siding. Axial plane crenulations with axial planes striking both  $100^{\circ}$  and  $180^{\circ}$  were observed within schists of the Peake Metamorphics and probably equate with the deformations producing the cross folds in the cover sediments.

Wopfner (1972) concluded that in the Permian there was syntectonic deposition, and fission track dating on a syenite sample from the Bungadillina Monzonite yielded an average age of  $266 \pm 23$  Ma (Radke, 1973).

Tertiary tectonism has caused uplift of huge fault blocks, forming the Peake and Denison Ranges. Jurassic and Cretaceous sediments and Tertiary silcretes are strongly upturned (dips to  $60^{\circ}$ ) adjacent to the ranges. Tectonism on the Mount Margaret Fault has been postulated as being late Miocene (Wopfner, 1972).

The sixth and final phase of tectonism is represented by the renewal of faulting during the Pleistocene along the Mount Margaret Fault which is indicated by a 150 m vertical displacement of the gypsite surface (Wopfner, 1968). Post-Cretaceous faulting and uplift along the Mount Margaret and Levi Faults has a combined throw in excess of 300 m (Wopfner, 1968).

#### METAMORPHISM

Previous comments on the grade of metamorphism for the Peake Metamorphics are restricted to a statement by Reyner (1955, p. 3) "Metamorphism of this group is due to granitisation as well as to regional processes".

From this present study three metamorphic episodes are evident;

1. amphibolite facies metamorphism of the Peake Metamorphics during the Carpentarian (approx. 1600 Ma),
2. ?greenschist facies metamorphism during the Musgravian Orogeny (approx. 1050 Ma),
3. low greenschist facies metamorphism of Peake Metamorphics and Adelaidean sediments during the Cambro-Ordovician Delamerian Orogeny (approx. 500 Ma).

Within the Peake Metamorphics two zones of different metamorphic grade have been recognised for the Carpentarian metamorphic event. Upper amphibolite facies metamorphism is evident from locally abundant migmatites northwest of the Wirriecurrie Granite. This is further supported by metamorphosed calcareous and pelitic sediments, in particular;

1. sillimanite gneisses and microcline-hornblende calc-silicates in the Coominaree Mine Block;

2. plagioclase-clinopyroxene granulites in the War Loan Mine Block;
3. plagioclase-microcline-epidote-hornblende calc-silicates in the Rockwater Hill Block;
4. clinopyroxene granulites at Lagoon Hill.

However, the good preservation of volcanic textures, and the mineralogy of rock types to the southwest and southeast of Peake ruins suggest only upper greenschist facies metamorphism in this area. Rock types are metavolcanics, quartz-muscovite schists, quartz-chlorite phyllites and micaceous metasandstones and quartzites.

Probable tectonism and low greenschist facies metamorphism during the Musgravian Orogeny are recorded by Rb-Sr and K-Ar geochronology on the Wirriecurrie Granite. Both Rb-Sr and K-Ar systems have been updated indicating a thermal event around 1050 Ma.

Metamorphism of basement rocks and Adelaidean sediments during the Delamerian Orogeny reached only low greenschist facies conditions. In general, the Adelaidean sediments exhibit little evidence of being metamorphosed. However, greenschist facies conditions are evident from rock types near Mount Young-husband and Mount Fox. Sandy dolomites occasionally exhibit porphyroblasts of potash feldspar, muscovite, chlorite and dolomite in a dolomite-quartz-biotite matrix, while metamorphosed shales exhibit lenticular quartz-rich aggregates in a lepidoblastic biotite-chlorite texture (Radke, 1976; 1977). A contact metamorphic aureole surrounds the largest Bungadillina Monzonite intrusive. Rock types in this zone include spotted, muscovite metasiltsstones, biotite-chlorite and biotite-muscovite meta-

siltstones, biotite metasandstones and quartz-biotite schists (Cooper, 1976; Stevenson 1976 b; Radke, 1977). Veins of quartz, feldspar, muscovite, chlorite and magnetite also occur within the contact aureole. Metamorphism during the Delamerian Orogeny was the last metamorphic episode; the rocks in the Peake and Denison Ranges have existed as a shallow or exposed basement block since the early Palaeozoic.

## ECONOMIC GEOLOGY

### HISTORICAL MINING ACTIVITIES

Alluvial gold was first discovered near Algebuckina in 1870 and near Peake Creek by the government geologist H.Y.L. Brown (1894). However, only a small amount of alluvial gold was found from numerous shallow pits on the north bank of the Neales River at Algebuckina.

During the late 1890's and early 1900's copper prospecting and mining flourished in the area. At that time there was a large community at the Peake overland telegraph station with 30 men employed in the mines. A smelter was built to treat copper ore from numerous mines and shafts and a water bore drilled near the telegraph station to a depth of 32.6 m, supplied 455 000 litres per day. Copper mining continued until 1920 and since then no mines have been worked. A description of the individual mines is presented below, and their locations are shown in Encl. 1.

#### 1. Copper Top Mines

Numerous shafts, pits and prospects occur in the vicinity of Peake ruins and near Copper Top Hill. These include Joes Prospect, Enrico Mine, Warrina Copper Mine, Coorunctite Mine, Central Mine, Warden and Denison, all of which are grouped under the Copper Top Mines because of the paucity of information.



In 1897 the Warrina Copper Mining Syndicate was formed, followed shortly afterwards in 1899 by the Amalgamated Copper Top Proprietary Company. The latter company erected a smelter, drilled a water bore and employed 30 men. Operations continued until 1904. By April, 1903, 242.8 tonnes of ore were treated and averaged 4% copper (Reyner, 1955). The deepest shaft was 35 m and the copper occurs as malachite, azurite and chalcocite in quartz-hematite veins (Reyner, 1955). The copper-bearing veins are located within basalts of the Tidnamurkuna Volcanics and Baltucoodna Quartzite and either strike parallel to the foliation or are cross-cutting striking northnorthwesterly parallel to the major faults.

## 2. Algebuckina Mine

The mine is located 1 km west of Algebuckina railway bridge in a sheared, gossanous zone within amphibolites, pegmatites and biotite-rich schists and gneisses. Two shafts were sunk, each 9 m deep. Little else is known of the early history of this mine. In 1970 Australasian Mining Corporation Limited drilled four diamond drill holes and the best intersection was 290 ppm  $U_3O_8$  between 24.00 and 26.21 m in DDH 1 (Sargeant, 1970).

## 3. Mount Kingston Prospect

The prospect lies 1.5 km north of Mount Kingston North within high grade metamorphics of the Peake Metamorphics, and is adjacent to a major north-south striking fault. The copper occurs as malachite and minor chalcocite in quartz veins.

## 4. Printa Mine

The Printa Mine is located 2 km north-northwest of Mount Kingston. It was first worked by the Pig Hill Copper Company in 1900. Surface shows were promising and 2 000 to 3 000 men were to

be employed, however only a series of small open cuts, two shafts (9.75 and 3.0 m deep) and an adit were dug. The copper mineralisation occurs within a small gossanous zone adjacent to pegmatites (Peake Metamorphics).

#### 5. Asbestos Mine

A malachite - crocidolite mine 2 km west of Tarlton Springs comprises several shallow pits within silty shales, dolomites and sandstones of the Duff Creek Beds. Malachite and blue, fibrous crocidolite are associated with quartz, biotite and ? goethite in hydrothermal aggregates. One specimen of dump material (6140 RS 15) contains greater than 1% Cu and 1% Mn (Encl. 4). A spherulitic potassium feldspar dyke outcrops at the mine and both the dyke and mineralisation are believed to be late-stage hydrothermal products from intrusion of the Bungadillina Monzonite. No early records of mining can be found.

#### 6. Last Chance Mine

The Last Chance Copper Mine is located 11.5 km south-southeast of Nilpinna Homestead and is on the margin of a large diapiric zone. It was worked from 1915 to 1919 under the name of W.J. and T.M. Briscoe. The main workings consisted of an open cut 18.3 x 6.1 x 8.2 m with an inclined shaft 11.6 m deep on one side of the open cut (Winton, 1919). The 0.5 - 0.6 m thick ore band, consisting of malachite, cuprite, chalcocite and chalcopyrite in a quartz-hematite lode, thinned with depth. Total ore shipments to 1918 were 162 tonnes containing 20-25% copper (Reyner, 1955).

#### 7. War Loan Mine

In 1916 two mineral claims were taken out by D. Clifford and J. Hillman who established a copper mine 8.5 km southeast of Nilpinna Homestead. An open cut 36.6 m x 4.3 m x 4.6 m

deep was excavated and about 6.1 m from the northern end an inclined shaft was sunk to a depth of 23.8 m, with several drives at the 12.2 m level (Winton, 1919). The claim was worked until 1920 and of a total of 149.5 tonnes of ore, copper contents ranged between 19 and 37%. Ore minerals were malachite and chalcocite, with traces of pyrite and gold in a ferruginous quartz lode (Reyner, 1955).

The mine is situated within blue-grey shales and siltstones of the War Loan Beds which tectonically overlie basalts of the Cadlareena Volcanics. Two diamond drill holes by North Broken Hill Limited (War Loan DDH 1 & 2) revealed several thin bands of anomalous copper values (Forwood, 1968; Western Mining Corporation, 1975).

#### 8. Coominaree Mine

The Coominaree Mine is located 6.5 km northeast of Nilpinna Homestead within diapiric, Adelaidean sediments. In 1916 W. Briscoe took out a mineral claim which became the Coominaree Mine. However only one shaft, 10.7 m deep and several shallow trenches and pits were sunk. Only 2 tonnes of copper ore are recorded to have been treated. The copper occurs as malachite in a quartz-calcite-hematite vein within disrupted dolomites and shales. The veins are associated with anomalous radioactivity and one specimen assayed 130 ppm  $U_3O_8$  (Iliff et al., 1974).

#### GEOCHEMISTRY

As part of the stratigraphic mapping programme, some of the main rock units were sampled and analysed to establish regional geochemical background values. A total of 83 specimens were analysed for up to 24 elements (Encl. 4). The values have been summarised in Table 12 which presents the average

COMPILED BY DRN I.M. CND		ROCK UNIT	ELEMENTS																										
			Ag	As	Au	Ba	Bi	Co	Cr	Cu	Li	Mn	Mo	Ni	P	Pb	Rb	Sb	Sn	Sr	Th	U	V	W	Zn	Zr	La	Y	
PEAKE AND DENISON RANGES MEAN ELEMENT ABUNDANCES FOR VARIOUS ROCK UNITS		BUNGADILLINA MONZONITE	x <sub>10</sub>	-	x <sub>12</sub>	1170 <sub>2</sub>	-	23 <sub>8</sub>	35 <sub>13</sub>	27 <sub>18</sub>	x <sub>2</sub>	-	x <sub>17</sub>	20 <sub>13</sub>	-	6 <sub>12</sub>	160 <sub>2</sub>	-	x <sub>15</sub>	1125 <sub>2</sub>	17 <sub>12</sub>	x <sub>12</sub>	101 <sub>12</sub>	x <sub>5</sub>	x <sub>12</sub>	115 <sub>2</sub>	-	-	
		CADLAREENA VOLCANICS	0 <sub>15</sub>	-	x <sub>12</sub>	922 <sub>7</sub>	x <sub>8</sub>	40 <sub>21</sub>	322 <sub>22</sub>	29 <sub>22</sub>	33 <sub>7</sub>	500 <sub>2</sub>	x <sub>7</sub>	95 <sub>22</sub>	-	54 <sub>22</sub>	81 <sub>7</sub>	-	-	142 <sub>7</sub>	x <sub>15</sub>	4 <sub>15</sub>	370 <sub>22</sub>	-	43 <sub>22</sub>	98 <sub>7</sub>	-	-	
		WIRRIE CURRIE GRANITE	-	-	x <sub>2</sub>	1225 <sub>2</sub>	-	50 <sub>2</sub>	17 <sub>2</sub>	62 <sub>2</sub>	33 <sub>2</sub>	-	x <sub>2</sub>	10 <sub>2</sub>	-	52 <sub>2</sub>	250 <sub>21</sub>	-	-	145 <sub>2</sub>	29 <sub>2</sub>	10 <sub>2</sub>	33 <sub>2</sub>	-	-	395 <sub>2</sub>	-	-	
		associated dykes	x <sub>3</sub>	x <sub>3</sub>	x <sub>3</sub>	767 <sub>3</sub>	x <sub>3</sub>	160 <sub>3</sub>	250 <sub>3</sub>	117 <sub>3</sub>	-	767 <sub>3</sub>	x <sub>3</sub>	217 <sub>3</sub>	400 <sub>3</sub>	5 <sub>3</sub>	-	x <sub>3</sub>	x <sub>3</sub>	-	-	-	333 <sub>3</sub>	x <sub>3</sub>	67 <sub>3</sub>	-	x <sub>3</sub>	50 <sub>3</sub>	
		Unnamed metamorphics Pd <sub>3</sub>	x <sub>4</sub>	x <sub>4</sub>	x <sub>4</sub>	367 <sub>4</sub>	x <sub>4</sub>	90 <sub>4</sub>	162 <sub>4</sub>	62 <sub>4</sub>	-	450 <sub>4</sub>	x <sub>4</sub>	137 <sub>4</sub>	675 <sub>4</sub>	2 <sub>4</sub>	-	x <sub>4</sub>	1 <sub>4</sub>	-	-	-	162 <sub>4</sub>	x <sub>4</sub>	20 <sub>4</sub>	-	x <sub>4</sub>	115 <sub>4</sub>	
		BALTUCOODNA QUARTZITE basalts	x <sub>4</sub>	x <sub>4</sub>	x <sub>5</sub>	300 <sub>5</sub>	x <sub>4</sub>	91 <sub>5</sub>	130 <sub>5</sub>	210 <sub>5</sub>	15 <sub>1</sub>	1450 <sub>4</sub>	x <sub>5</sub>	156 <sub>5</sub>	550 <sub>4</sub>	7 <sub>5</sub>	50 <sub>1</sub>	x <sub>4</sub>	x <sub>4</sub>	95 <sub>1</sub>	x <sub>1</sub>	14 <sub>1</sub>	314 <sub>5</sub>	x <sub>4</sub>	78 <sub>5</sub>	165 <sub>1</sub>	x <sub>4</sub>	92 <sub>4</sub>	
		metasediments	x <sub>6</sub>	x <sub>6</sub>	x <sub>6</sub>	717 <sub>6</sub>	x <sub>6</sub>	73 <sub>6</sub>	138 <sub>6</sub>	50 <sub>6</sub>	-	562 <sub>6</sub>	x <sub>6</sub>	125 <sub>6</sub>	550 <sub>6</sub>	7 <sub>6</sub>	-	x <sub>6</sub>	1 <sub>6</sub>	-	-	-	200 <sub>6</sub>	x <sub>6</sub>	50 <sub>6</sub>	-	92 <sub>6</sub>	70 <sub>6</sub>	
		Unnamed schists Pd <sub>2</sub>	x <sub>2</sub>	x <sub>2</sub>	x <sub>2</sub>	500 <sub>2</sub>	x <sub>2</sub>	125 <sub>2</sub>	200 <sub>2</sub>	75 <sub>2</sub>	-	500 <sub>2</sub>	10 <sub>2</sub>	200 <sub>2</sub>	750 <sub>2</sub>	2 <sub>2</sub>	-	x <sub>2</sub>	x <sub>2</sub>	-	-	-	200 <sub>2</sub>	x <sub>2</sub>	25 <sub>2</sub>	-	x <sub>2</sub>	45 <sub>2</sub>	
		TIDNAMURKUNA VOLCANICS rhyolites	x <sub>2</sub>	x <sub>2</sub>	x <sub>4</sub>	740 <sub>4</sub>	x <sub>2</sub>	58 <sub>4</sub>	44 <sub>4</sub>	90 <sub>4</sub>	12 <sub>2</sub>	550 <sub>2</sub>	x <sub>4</sub>	51 <sub>4</sub>	500 <sub>2</sub>	22 <sub>4</sub>	235 <sub>2</sub>	x <sub>2</sub>	1 <sub>2</sub>	55 <sub>2</sub>	32 <sub>2</sub>	10 <sub>2</sub>	72 <sub>4</sub>	x <sub>2</sub>	36 <sub>4</sub>	405 <sub>2</sub>	75 <sub>2</sub>	50 <sub>2</sub>	
basalts	x <sub>3</sub>	x <sub>1</sub>	x <sub>4</sub>	700 <sub>4</sub>	x <sub>3</sub>	151 <sub>4</sub>	222 <sub>4</sub>	185 <sub>4</sub>	25 <sub>1</sub>	933 <sub>3</sub>	x <sub>4</sub>	237 <sub>4</sub>	666 <sub>3</sub>	6 <sub>4</sub>	95 <sub>1</sub>	x <sub>3</sub>	x <sub>3</sub>	100 <sub>1</sub>	8 <sub>1</sub>	x <sub>1</sub>	72 <sub>4</sub>	x <sub>3</sub>	54 <sub>4</sub>	165 <sub>1</sub>	x <sub>3</sub>	73 <sub>3</sub>			
Unnamed metamorphics Pd <sub>1</sub>	x <sub>6</sub>	x <sub>6</sub>	x <sub>6</sub>	533 <sub>6</sub>	x <sub>6</sub>	110 <sub>6</sub>	212 <sub>6</sub>	118 <sub>6</sub>	-	700 <sub>6</sub>	x <sub>6</sub>	185 <sub>6</sub>	1380 <sub>6</sub>	5 <sub>6</sub>	-	x <sub>6</sub>	x <sub>6</sub>	-	-	-	208 <sub>6</sub>	x <sub>6</sub>	46 <sub>6</sub>	-	50 <sub>6</sub>	93 <sub>6</sub>			
SCALE		<div><div>115<sub>22</sub></div><div>x<sub>6</sub></div><div>-</div></div> <div>Values in p.p.m., number of analyses shown as subscripts.</div> <div>Analysed but not detected, or average falls below detection limit.</div> <div>Not analysed.</div>																											
		The averages are derived from values quoted in enclosure 4.																											

TABLE 12

82A.

88

The averages are derived from values quoted in enclosure 4.

TABLE 12

DEPARTMENT OF MINES AND ENERGY  
SOUTH AUSTRALIA

SCALE

DATE 3-4-78

PLAN NUMBER

S13334

elemental abundances for the Bungadillina Monzonite, Cadlarena Volcanics, Wirriecurrie Granite and all units within the Peake Metamorphics. However a comparison of the results obtained with the elemental abundances of the respective rock types (Turekian & Wedepohl, 1961; Hawkes & Webb, 1962) reveal no significantly anomalous values in this regional rock sampling programme.

Analyses of basalts within the Peake Metamorphics generally show higher elemental abundances than other rock types, but the values are typical for basaltic rocks (Hawkes & Webb, 1962). Hydrothermal quartz-hematite-malachite veins are common within the basalts, particularly near the upper contact between basalts and overlying metasediments. Many of the old mines are located in this environment, e.g. the shafts near Peake ruins and Mount Denison. The basalt - metasediment association provides a good target for future base metal exploration in the Peake and Denison Ranges.

#### SUMMARY

Detailed mapping of the pre-Mesozoic rocks of the Peake and Denison Ranges has enabled a revised stratigraphic succession to be established. The ? Lower Proterozoic Peake Metamorphics have been subdivided into six units while the unconformably overlying Adelaidean sediments have been subdivided into twenty one units.

The Peake Metamorphics comprise a minimum of 11 500 m of dominantly volcanics and coarse-grained arenaceous sediments. The volcanics are chiefly porphyritic and amygdaloidal basalts with minor flow-banded, porphyritic rhyolites.

Greenschist to amphibolite facies metamorphism and deformation of the Peake Metamorphics, and syntectonic granite intrusion (Wirriecurrie Granite) has been dated at about 1600 Ma.

The updating of Rb-Sr and K-Ar systems, revealed by the geochronological studies, indicate probable mild tectonism and greenschist facies metamorphism about 1050 Ma (Musgravian Orogeny).

Overlying the Peake Metamorphics are at least 29 000 m of Adelaidean sediments. Deposition occurred in a steadily subsiding, fault-bounded trough with shallow-water sediments predominating. Clastic sediments were chiefly derived from the Gawler Craton to the west and southwest.

Adelaidean sedimentation is first recorded by the Willouran Callanna Beds. The basal unit of conglomerates and quartzites (Younghusband Conglomerate) is overlain by stromatolitic dolomites (Coominaree Dolomite) and vesicular basalts and altered dolerites of the Cadlareena Volcanics. These three units correlate with a similar basal Adelaidean succession in the Mount Painter Province and this correlation is further supported by the major and trace element chemistry of the volcanics in the two areas.

Diapirism has caused extensive disruption of the sequence immediately above the Cadlareena Volcanics. From the dismembered succession the following stratigraphic units (of Willouran or Torrensian age) have been established, Rockwater Beds, War Loan Beds, Nilpinna Beds, Duff Creek Beds, and Murrana Beds. Lithologies and the abundance of mud cracks, ripple marks, clay galls, salt and gypsum casts indicate a paralic environment of deposition.

Sequences of Torrensian age which can be definitely assigned to the Burra Group include; unnamed siltstone unit, Fountain Spring Beds, Mount Margaret Quartzite, Skillogalee Dolomite, unnamed transition unit and Kalachalpa Formation. Deposition during the Torrensian occurred in a paralic environment as indicated by the presence of intraformational dolomitic and magnesitic conglomerates, stromatolitic dolomites and oolitic sediments. Over 10 000 m of Burra Group sediments outcrop in the Peake and Denison Ranges, which

is a thicker and more arenaceous sequence than elsewhere in the Adelaide Geosyncline. For example in the Mount Painter Province (Coats and Blisset, 1971) Willouran and Torrensian sediments total 6 500 m compared with 27 000 m in the Peake and Denison Ranges thus indicating a prolonged period of more rapid basinal subsidence and sedimentation in the Ranges.

A period of mild erosion preceded deposition of the Sturtian Umberatana Group. The onset of glacial conditions resulted in deposition of the glacio-marine Calthorinna Tillite and interfingering sandstones. The overlying marine siltstones and shales of the Tapley Hill Formation are, in turn, overlain by a regressive sequence of dolomites (Thora Dolomite), and fine sandstones and reddish silty shales (unnamed siltstone of the Willochra Subgroup).

Folding, faulting and diapirism of the Adelaidean succession during the Cambro-Ordovician Delamerian Orogeny was accompanied by low greenschist facies metamorphism and intrusion of quartz-poor porphyritic monzonites, syenites, diorites and albitites (Bungadillina Monzonite).

Outcropping Permian gritty limestones and diamictites have been assigned to the Boorthanna Formation. Clast lithologies within the diamictites are dominantly of local Adelaidean rock types, and striated, faceted and soled clasts indicate a glacial origin.

Analyses of 83 specimens (for up to 24 elements) revealed no significant anomalous values in a regional rock sampling program. However the basalt - metasediment association within the Peake Metamorphics provides a good target for future base metal exploration in the Peake and Denison Ranges.

### ACKNOWLEDGEMENTS

Special thanks are due to Mr. R.P. Coats and Dr. W.V. Preiss for their aid and discussions during the preparation of this paper and, along with Mr. B.P. Thomson, C.R. Dalgarno Dr. B.G. Forbes for critically reading the manuscript. We acknowledge Dr. T.R. Fairchild for permission to use Fig. 20 from his Ph.D thesis and R. Wells for documenting details on mines and previous mining activities.

We also express our appreciation for the co-operation and hospitality of the managers of the pastoral properties in the area.



# REFERENCES

- Allchurch, P.D., Wopfner, H., Harris, W.K. and McGowran, B.,  
1973. Cootanoorinna No. 1 Well. Rep. Invest., geol. Surv.  
S. Aust., 40: 80 pp.
- Badiozamani, K., 1973. The Dorag dolomitization model - application  
to the Middle Ordovician of Wisconsin. J. sedim. Petrol.  
43: 965-984.
- Barclay, C.J., 1974. Relinquishment report E.L. 108; Shell  
Development (Aust) Pty. Ltd., S. Aust. Dept. Mines  
open file Env. 2388 (unpublished).
- Benbow, M.C., and Flint, R.B., 1978. The Engenina Adamellite and  
Balta Granite of the Mount Woods Inlier. S. Aust.  
Dept. Mines and Energy report (unpublished).
- Brown, H.Y.L., 1894. Report on the Peake and Denison Ranges and  
adjoining country with special reference to the occurrence  
of gold. Parl. Pap. S. Aust. No. 25.
- Carey, S.W. and Ahmad, N., 1960. Glacial marine sedimentation.  
In: Geology of the Arctic. Raasch, G.O. Toronto, Univ.  
Toronto Press, pp. 865-894.
- Chugg, R.I., 1957. The hydrogeology of a portion of the Great  
Artesian Basin near the Peake and Denison Ranges. Rep.  
Invest., geol. Surv. S. Aust., 10: 27 pp.
- Coats, R.P., 1964a. The geology and mineralisation of the Blinman  
Dome Diapir, Rep. Invest., geol. Surv. S. Aust., 6: 23 pp.
- Coats, R.P., 1964b. Umberatana Group. Q. geol. Notes, geol. Surv.  
S. Aust., 9: 7-12.
- Coats, R.P. and Blisset, A.H., 1971. Regional and economic geology  
of the Mount Painter Province. Bull. geol. Surv. S. Aust.,  
43: 426pp.
- Coats, R.P., 1973. COPLEY, South Australia. Explanatory Notes,  
1:250 000 geological series. Sheet Sh/54-9. Geol. Surv.  
S. Aust.

- Coats, R.P. and Forbes, B.G., 1977. Evidence for two Sturtian Glaciations in South Australia - a reply. Q. geol. Notes, geol. Surv. S. Aust., 64: 19-20.
- Cooper, P.F. and Tuckwell, K., 1971. The Upper Precambrian Adelaidean of the Barrier Ranges, a new subdivision. Q. geol. Notes, geol. Surv. N.S.W., 3: 8-16.
- Cooper, R., 1976. Petrographic description of twenty five samples from the Peake and Denison Ranges. Amdel report MP365/77 (unpublished).
- Cooper, R., Couppey, C., Jacque, M., Leslie, W. and Magnier, P., 1963. Mission geologique sur les bordures occidentales du Grand Bassin Artesian, French Petroleum Company. S. Aust. Dept. Mines open file Env. 386/2 (unpublished).
- Crawford, A.R., 1963. The Wooltana Volcanic belt, South Australia. Trans. R. Soc. S. Aust., 87: 123-154.
- Dalgarno, C.R. and Johnson, J.E. 1968. Diapiric Structures in Flinders Ranges, South Australia. In: Braunstein, J. and O'Brien, G.D. (Eds), Diapirism and Diapirs. Mem. Am. Ass. Petrol. Geol. 8: 301-314.
- Derrick, G.M., Wilson, I.H., and Hill, R.M., 1976. Revision of stratigraphic nomenclature in the Precambrian of north-western Queensland. Qld. Govt. Mining J. 77: 300-306.
- Derrick, G.M., Wilson, I.H., Hill, R.M., Glikson, A.Y. and Mitchell, J.E., 1977. Geology of the Mary Kathleen 1:100 000 sheet area, northwest Queensland. Bull. Bur. Miner. Resour. Geol. Geophys. Aust., 193: 114 pp.
- Dickinson, S.B., Parkin, L.W., Pitman, R.K., Reyner, M.L. and Hughes F.E., 1954a. Algebuckina map sheet, Geological Atlas of South Australia, 1:63 000 series, Geol. Surv. S. Aust.

- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K. 1954b. Nilpinna map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K., 1955a. Anna map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K., 1955b. Boorthanna map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K., 1955c. Umbum map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K., 1955d. Conway map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- Dickinson, S.B., Parkin, L.W., Hughes, F.E., Reyner, M.L. and Pitman, R.K., 1955e. Cadlareena map sheet, Geological Atlas of South Australia, 1:63 000 series. Geol. Surv. S. Aust.
- East, J.J., 1889. On the geological structure and physical features of central Australia. Trans. R. Soc. S. Aust., 12: 31-53.
- Ewart, A., 1971. Chemical changes accompanying spherulitic crystallization in rhyolitic lavas, Central Volcanic Region, New Zealand. Mineralog. Mag. 38: 424-434.
- Fairchild, T.R., 1975. The geological setting and palaeobiology of a late Precambrian stromatolitic microflora from South Australia. PhD. thesis, University of California, 271 pp (unpublished).
- Fander, H.W., 1963. The Wooltana Lavas Trans. R. Soc. S. Aust., 87: 155-157.

- Forbes, B.G., 1961. Magnesite of the Adelaide System: a discussion of its origin. Trans. R. Soc. S. Aust., 85: 217-222.
- Forbes, B.G., 1964. The River Wakefield Group west of Clare. Q. geol. Notes, geol. Surv. S. Aust., 11: 6-7.
- Forbes, B.G., 1974. Bungaree Quartzite (new name): Lower Adelaidean, southwest of Spalding, South Australia. Q. geol. Notes, geol. Surv. S. Aust., 63: 9-14.
- Forwood, P.S., 1968. Final report on exploration on Special Mining Leases 125 and 134, Peake and Denison Ranges, South Australia. S. Aust. Dept. Mines open file Env. 2072 (unpublished).
- Freytag, I.B., 1966. Proposed rock units for marine lower Cretaceous sediments in the Oodnadatta region of the Great Artesian Basin. Q. geol. Notes, geol. Surv. S. Aust., 18: 3-7.
- Freytag, I.B., Heath, G.R., and Wopfner, H., 1967. OODNADATTA map sheet, Geological Atlas of South Australia, 1:250 000 series. Geol. Surv. S. Aust.
- Hawkes, H.E. and Webb, J.S., 1962. Geochemistry in Mineral Exploration. Harper and Row, New York, 415 pp.
- Heath, G.R., 1965. Permian sediments of the Mount Dutton Inlier. Q. geol. Notes, geol. Surv. S. Aust., 14: 3-5.
- Holmes, D., 1970. South Australia Potash Project, Special Mining Lease 329, Occidental Mins Corp. (Aust.). S. Aust. Dept. Mines open file Env. 1313 (unpublished).
- Holmes, D.A. and Rayment, P., 1970. Oxymin Boorthanna No. 1 Well completion report. Occidental Mins Corp. (Aust.). S. Aust. Dept. Mines open file Env. 1241 (unpublished).
- Iliff, G., 1975. Exploration 1974 on E.L. 100, Peake-Denison and Mt. Margaret Ranges, South Australia. Uranerz (Aust.) Pty Ltd, S. Aust. Dept. Mines open file Env. 2381 (unpublished).

- Iliff, G.D., Robinson, P. and Johnson, J., 1974. Exploration 1973 on E.L. 33 Mt. Margaret Range, Peake and Denison Block, South Australia. Uranerz (Aust.) Pty Ltd , S. Aust. Dept. Mines open file Env. 2253 (unpublished).
- Jack, R.L., 1930. Geological structure and other factors in relation to underground water supply in portions of South Australia. Bull. geol. Surv. S. Aust., 14: 48 pp.
- Laut, P., Keig, G., Lazarides, M., Loffler, E., Margules, C., Scott, R.M. and Sullivan, M.E., 1977. Environment of South Australia - Province 8 Northern Arid. Division of Land Use Research, Commonwealth Scientific and Industrial Research Organisation, Canberra Australia, pp. 152-155.
- Leeson, B., 1972. Geochemical and geophysical investigations near the Last Chance Mine. Mineral Resour. Rev., S. Aust., 132: 95-102.
- Le Maitre, R.W., 1976. The chemical variability of some common igneous rocks. J. Petrology 17 (4): 589-637.
- Ludbrook, N.H., 1961. Permian to Cretaceous subsurface stratigraphy between Lake Phillipson and the Peake and Denison Ranges, South Australia. Trans. R. Soc. S. Aust., 85: 67-80.
- Mawson, D., 1927. Geological notes on an area along the northwestern margin of the Willouran Range. Trans. R. Soc. S. Aust., 51: 386-390.
- Mawson, D. and Sprigg, R.C., 1950. Subdivision of the Adelaide System. Aust. J. Sci., 13 (3): 69-72
- Mirams, R.C. and Forbes, B.G., 1964. Burra Group. Q. geol. Notes, geol. Surv. S. Aust., 9: 5-7.
- Murrell, B., 1977. The stratigraphy and tectonics across the Torrens Hinge Zone <sup>between Andamooka and Marree, South Australia.</sup> / Ph.D. thesis, University of Adelaide (unpublished).

- Papalia, N., 1970. Final report on Weedina No. 1 Well. Pexa Oil N.L., S. Aust. Dept. Mines open file Env. 1374 (unpublished).
- Pearce, J.A., 1976. Statistical analysis of major element patterns in basalts. J. Petrology 17: 15-43.
- Preiss, W.V., 1973a. Early Willouran stromatolites from the Peake and Denison Ranges and their stratigraphic significance. S. Aust. Dept. Mines report 73/208 (unpublished).
- Preiss, W.V., 1973b. Palaeoecological Interpretations of South Australian Precambrian Stromatolites. J. geol. Soc. Aust., 19: 501-509.
- Preiss, W.V., 1974. The River Broughton Beds - A Willouran sequence in the Spalding Inlier. Q. geol. Notes, geol. Surv.S. Aust., 49: 2-8.
- Radke, F., 1973. Fission - track dating. Amdel report Project 1/1/148, Progress report No. 1 (unpublished).
- Radke, F., 1976. Examination of rocks from WARRINA 1:250 000 sheet. Amdel report MP 1580/76 (unpublished).
- Radke, F., 1977. Petrography of thirteen rocks from the WARRINA 4 mile sheet, Peake and Denison Ranges. Amdel report MP1648/77 (unpublished).
- Reyner, M.L., 1955. The geology of the Peake and Denison Region. Rep. Invest., geol. Surv. S. Aust., 6: 23 pp.
- Sargeant, F.J., 1970. Report on exploration during the period 15/2/70 to 15/8/70, S.M.L. 270, South Australia. Australasian Mining Corporation Ltd, S. Aust. Dept. Mines open file Env. 1015 (unpublished).
- Schermerhorn, L.J.G., 1966. Terminology of mixed coarse-fine sediments. J. Sedim. Petrol., 36: 834 pp.
- Schopf, J.W. and Fairchild, T.R., 1973. Late Precambrian micro-fossils: a new stromatolitic biota from South Australia. Nature 242: 537-538.

- Scoular, G., 1887. Sketch of the geology of the southern and western parts of the Lake Eyre Basin. Trans. R. Soc. S. Aust., 9: 39-54.
- Smith, N., 1973. Gravity and magnetic analysis, Lagoon Hill Area, South Australia. Company report E.L. 22; Chevron Exploration Corporation. S. Aust. Dept. Mines open file Env. 2182 (unpublished).
- Spencer, A.M., 1971. Late Precambrian glaciation in Scotland. Mem.geol. Soc. Lond., 6: 98 pp.
- Sprigg, R.C., 1949. Thrust structures in the Witchelina area, South Australia. Trans. R. Soc. S. Aust., 73 (1): 40-47.
- Sprigg, R.C., 1952. Sedimentation in the Adelaide Geosyncline and the formation of the continental terrace. Sir Douglas Mawson Anniv. Vol. Univ. of Adelaide, pp 153-159.
- Steveson, B.G., 1973. The geochronology of the eastern basement rocks. Amdel report project 1/1/140; progress report No. 2 (unpublished).
- Steveson, B.G., 1976a. Examination of rocks from the Cadlareena Volcanics. Amdel report MP 224/77 (unpublished).
- Steveson, B.G., 1976b. Examination of eleven rocks from WARRINA 1:250 000 sheet. Amdel report MP 335/77 (unpublished).
- Steveson, B.G., 1976c. Examination of rocks from WARRINA 1:250 000 sheet. Amdel report MP 1553/76 (unpublished).
- Stuart, J.M., 1860. J.M. Stuart's exploration in 1860. Parl. Pap. S. Aust., No. 65.
- Stuart, J.M., 1862. J.M. Stuart's exploration, 1861-1862. Parl. Pap. S. Aust., No. 21.
- Teluk, J.A., 1974. Completion report on E.L. 22, Lagoon Hill Area - South Australia. Chevron Exploration Corporation. S. Aust. Dept. Mines open file Env. 2182 (unpublished).

- Thomson, B.P. and Coats, R.P., 1964. The Callanna Beds. Q. geol. Notes, geol. Surv. S. Aust., 9: 3-5.
- Thomson, B.P., 1966. The lower boundary of the Adelaide System and older basement relationships in South Australia. J. geol. Soc. Aust., 13, (1): 203-213.
- Thomson, B.P., 1969. The Adelaide System. In: Parkin, L.W., (Ed.). Handbook of South Australian Geology. Geol. Surv. S. Aust., Gov. Printer, Adelaide, pp 69-71.
- Thomson, B.P., 1974. Tectonics and regional geology of Willyama, Mount Painter and Denison Inlier areas. In: Economic geology of Australia and Papua New Guinea - I. Metals. Aust. Inst. Min. and Met. Monograph Series, 5, pp 469-475.
- Townsend, I.J., 1973. A synthesis of stratigraphic drilling of the Arckaringa Basin 1969-1971. S. Aust. Dept. Mines report 73/87 (unpublished).
- Townsend, I.J. and Ludbrook, N.H., 1975. Revision of Permian and Devonian nomenclature of four formations in and below the Arckaringa Basin. Q. geol. Notes, geol. Surv. S. Aust., 54: 1-5.
- Turekian, K.K. and Wedepohl, K.H., 1961. Distribution of the elements in some major units of the earth's crust. Bull. geol. Soc. Am., 72: 175-192.
- Turner, A.R., 1969. Some aspects of the basic lavas of the Gawler Platform. Amdel report MP 809/69 (unpublished).
- Turner, A.R., 1969. The geochronology of stratigraphically significant rocks from South Australia. Amdel report Project 1/1/126, Progress report No. 2 (unpublished).
- Vennum, W.R. and Eberlein, G.D., 1977. Spherulitic rhyolite dyke from Goat Island, southeastern Alaska. J. Research U.S. geol. Surv., 5 (4): 445-451.



- Webb, A.W., 1970. The geochronology of stratigraphically significant rocks from South Australia. Amdel report Project 1/1/126, Progress report No. 4 (unpublished).
- Webb, A.W., 1974. The geochronology of the eastern basement rocks. Amdel report Project 1/1/140, Progress report No. 3 (unpublished).
- Webb, A.W., 1976. Geochronology of the granitic rocks of southeastern South Australia. Amdel final report Project 1/1/128 (unpublished).
- Webb, A.W., 1977a. Geochronology of stratigraphically significant rocks from South Australia. Amdel report Project 1/1/126, Progress report No. 16 (unpublished).
- Webb, A.W., 1977b. Geochronology of sample P 266/76. Amdel report Project 1/1/126, Progress report No. 15 (unpublished).
- Webb, A.W., and Lowder, G.G., 1971. The geochronology of stratigraphically significant rocks from South Australia. Amdel report project 1/1/126, Progress Report No. 5 (unpublished).
- Webb, A.W. and Lowder, G.G., 1972. The geochronology of the eastern basement rocks. Amdel report Project 1/1/140, Progress report No. 1 (unpublished).
- Western Mining Corporation, 1975. Exploration Licence 192, Peake and Denison Ranges. S. Aust. Dept. Mines open file Env. 2525 (unpublished).
- Whitehead, S. 1972a. Metamorphic and sedimentary rocks from Mt. Kingston. Amdel report MP 1915/72 (unpublished).
- Whitehead, S., 1972b. Petrographic description of eleven rock specimens from WARRINA 1:250 000 sheet. Amdel report MP 2151/72 (unpublished).

- Williams, G.E. 1975. Late Precambrian glacial climate and the Earth's obliquity. Geol. Mag., 112 (5): 441-465.
- Wilson, J.L., 1975. Carbonate Facies in Geological History. Springer - Verlag, New York, 471 pp.
- Winton, L.J., 1919. Report on the Last Chance Copper Mine and War Loan Copper Mine. Min. Rev., Adelaide, 29: 54-56.
- Wopfner, H., 1968. Cretaceous sediments on the Mount Margaret Plateau and evidence for neo-tectonism. Q. geol. Notes, geol. Surv. S. Aust., 28: 7-11.
- Wopfner, H. 1970. Permian palaeogeography and depositional environment of the Arckaringa Basin, South Australia. In: Second Gondwana Symposium Natal, Witwatersrand press, Natal, pp 273-291.
- Wopfner, H., 1972. Depositional history and tectonics of South Australian sedimentary basins. Mineral Resour. Rev., S. Aust., 133: 32-50.
- Wopfner, H., Freytag, I.B. and Heath, G.R., 1970. Basal Jurassic - Cretaceous rocks of the western Great Artesian Basin, South Australia: Stratigraphy and environment. Bull. Am. Ass. Petrol. Geol., 54: 383-416.
- Wopfner, H. and Heath, G.R., 1963. New observations on the basal Cretaceous-Jurassic sandstone in the Mt. Anna Region, South Australia. Aust. J. Sci. 26: 57-59.
- Young, G.M., 1976. Iron-formation and glaciogenic rocks of the Rapitan Group, Northwest Territories, Canada. Precambrian Res. 3: 137-158.
- Youngs, B.C., 1975a. The geology and hydrocarbon potential of the Pedirka Basin. Rep. Invest., geol. Surv. S. Aust., 44: 47 pp.
- Youngs, B.C., 1975b. The Early Permian Purni Formation of the Pedirka Basin. Q. geol Notes, geol. Surv. S. Aust., 54: 5-12.

*				
RB 79/19 10A				
AGE	STRATIGRAPHIC UNIT: SYMBOL AND PRIMARY REFERENCES	THICKNESS (metres)	LITHOLOGY	REMARKS
ORDOVICIAN	Bungadillina Monzonite (Odb) Reyner, 1955	-	Coarse-grained porphyritic monzonites and syenites, porphyritic diorites, porphyritic white albitites.	Intrusives are plagioclase rich and quartz poor.
	ADELAIDEAN STRATIGRAPHY			
MARTINIAN	Unnamed siltstone (Ph1) Thomson, in Parkin, 1969	950 (minimum)	Red-brown silty shales, grey-green shales, silty dolomites, fine sandstones and thin grey-yellow dolomites.	Records the first appearance of reddish siltstones in the Umeratana Group-hence allocated to the Willochra Subgroup.
	*Thora Dolomite (Pfl) Mawson and Sprigg, 1950	30-40	Buff weathering, grey-green and brown dolomites, minor siltstones; large festoon surfaces and algal bedding.	Characterized by a light coloured weathering pattern which distinguishes it from adjacent units.
	Tapley Hill Formation (Eft) Coats, 1964; 1971	150	Laminated grey-green silty shales and silty dolomites; basal thin sandy dolomite.	Exhibits characteristic fine lamination; records a marine transgression following the pre-existing glacial conditions.
STURTIAN	Unnamed sandstone (Euf)	530	Quartzitic sandstones, arkoses, argillaceous sandstones; red porphyry granules and large scale cross-bedding.	Fluvio-glacial; intertongues with the Calthorinna Tillite and marks the end of the glacial period.
	*Calthorinna Tillite (Eub) Coats, 1964; 1971 Coats and Forbes, 1977 Reyner, 1955	650 (minimum)	Diamictites, conglomeratic dolomites, pale green shales, gritty sandstones, arkoses and quartzites.	Represents the upper phase of lower Sturtian glacial activity. Interbedded diamictites (glacial) and marine sediments indicate a glacial marine environment.
DISCONFORMITY				
TORENSIAN	*Kalachalpa Formation (Pbk) Fairchild, 1975 Coats, 1973	900	Grey-green and brown siltstones and shales, gritty sandstones and quartzites, conglomeratic dolomites, stromatolitic dolomites, oolitic sediments, black cherts; quartzites and shales at the top.	Stromatolitic microflora described by Fairchild, 1975; considered to be equivalent to the Myrtle Springs Formation; gradational contact with the underlying unnamed unit.
	Unnamed unit (Eb2) Fairchild, 1975	2000 (minimum)	Gritty quartzitic sandstone, stromatolitic dolomites, conglomeratic dolomites, magnesite conglomerates, black and minor red cherts, grey shales and siltstones; mud cracks ripple marks, cross-bedding.	Many lithological affinities with the Skillogalee Dolomite e.g. dolomite pebble conglomerates, magnesites and black cherts; this unit is probably transitional between the Kalachalpa Formation and Skillogalee Dolomite.
	GAP IN THE SEDIMENTARY RECORD (gradational contact suspected)			
	Skillogalee Dolomite (Ebk) Mirams and Forbes, 1964. Reyner, 1955	3600 (minimum)	Basal member composed of quartzites, sandstones, shales and minor dolomites. The middle member consists of sandstones, dark and pale grey conglomeratic dolomites, magnesite conglomerates and blue-black cherts. The upper member is mainly dark grey conglomeratic dolomite.	Outcrops in a series of basal synclines forming the core of Margaret Inlier; Stromatolite type <u>Baicalia burra</u> is common in the upper two members; lower contact with the Mount Margaret Quartzite is conformable although often disrupted by diapirism.
TORENSIAN	*Mount Margaret Quartzite (Pbo) Reyner, 1955 Thomson and Coats, 1964	2500	White orthoquartzites, slaty quartzitic sandstones, dark grey sandy siltstones, green-grey silty shales, minor dolomitic siltstones near the base; thick orthoquartzites at the top; clay galls, ripple marks and cross-bedding.	Comprises Mount Margaret and also constitutes the Mount Margaret Plateau. The lower contact with the Fountain Spring Beds is gradational, the passage downwards being marked by an increased siltstone content.
	*Fountain Spring Beds (Ebl) Forbes, 1964	1100 (minimum)	Characteristic laminated grey dolomitic siltstones, interbedded grey quartzites with clay gall laminations, grey-green silty shales containing salt casts, minor grey silty dolomites.	The upper contact with the Mount Margaret Quartzite is marked by an increase in the frequency of orthoquartzites; laminated dolomitic siltstones and silty dolomites are similar to lithologies in the underlying unnamed siltstone; considered to be River Wakefield Subgroup equivalent.
	Sedimentary contact not observed-gradational contact suspected on the basis of lithological overlaps			
	Unnamed siltstone (Eb1)	1200 (minimum)	Laminated grey pyritic dolomitic siltstones, green shales, pyritic silty dolomites, fine sandstones and minor quartzites; ripple marks, cross-beds, salt casts.	Dark grey laminated siltstones are typical of the Fountain Spring Beds; some of the dolomites and sandstones (salt casts) are reminiscent of the Duff Creek Beds.
WILLOUREAN OR TORENSIAN	GAP IN THE SEDIMENTARY RECORD			
	Undifferentiated blocks in diapirs and faulted sequences (Ea)	5000 (minimum)	Lithologies from 8 blocks are described in the text (see Page ). Sediments observed in diapirs often contain an abundance of salt casts.	Most of the sediments in these blocks are derived from lower Adelaidean sequences, above the Cadlareena Volcanics, which have been dismembered by diapirism and faulting.
	GAP IN THE SEDIMENTARY RECORD			
	*Murrana Beds (Eaa)	2900 (minimum)	Laminated gritty quartzitic sandstones, grey-green silty shale interbeds; arkoses, pebbly dolomites, purple and grey silty shales near the top; ripple marks, mud cracks, cross-beds, flute casts and salt casts.	This unit has lithological affinities with the Duff Creek Beds e.g. minor pale dolomites in the lower part of the unit and fine sandstones and green shales both containing salt casts.
WILLOUREAN OR TORENSIAN	Sedimentary contact not observed-gradational contact suspected on the basis of lithological overlaps			
	*Duff Creek Beds (Eaf) Reyner, 1955 Thomson and Coats, 1964	5500 (minimum)	Laminated olive green silty shales and thin arkose interbeds, flaggy grey and yellow dolomites, dolomitic siltstones, pyritic fine sandstones, minor quartzites; mud cracks, ripple marks, clay galls, cross-bedding; horizons with abundant salt and gypsum casts.	Pale buff, flaggy dolomites characterise the unit; deposited in lagoonal, supratidal and intertidal environments; fine platy sandstones (salt casts) in the lower part of the sequence are similar to the Nilpinna Beds.
	Sedimentary contact not observed-gradational contact suspected on the basis of lithological overlaps			
	*Nilpinna Beds (Ean)	2100 (minimum)	Fine sandstones, grey-green silty shales, quartzites and minor grey silty dolomites and green arkoses; mud cracks, cross-bedding and salt casts.	Blue-grey shales and fine-medium sandstones (festoon cross-beds, salt casts) near the base of the unit are similar to the War Loan Beds.
WILLOUREAN	Sedimentary contact not observed-gradational contact suspected on the basis of lithological overlaps			
	*War Loan Beds (Eaw)	600 (minimum)	Blue-grey shales and siltstones, dark grey silty dolomites and greenish feldspathic sandstones and arkoses.	Blue-grey colour of the sediments is characteristic.
	GAP IN THE SEDIMENTARY RECORD			
	*Rockwater Beds (Ead)	200 (minimum)	Blue-grey and black cherts (weather to a cream colour), grey-black pebbly dolomites, black shales and siltstones and quartzitic sandstones.	This unit outcrops only in small blocks stratigraphically isolated by faulting and diapirism; the sequence probably lies in close proximity, stratigraphically, to Cadlareena Volcanics.
WILLOUREAN	GAP IN THE SEDIMENTARY RECORD - sequence dismembered by diapirism & faulting			
	*Cadlareena Volcanics (Eoc) Coats and Blissett, 1971 Thomson, 1966 Preiss, 1973	750 (minimum)	Vesicular basalts and altered dolerites; minor andesites, dacites, and rhyolites; tuffs and lapilli tuffs; minor lenticular reddish mudstones and quartzites with red shale interbeds near the base and top of the unit.	The Volcanics mark the last recognised unit of the Callanna Beds; equivalents are recognised in the Willouran Ranges and Mount Painter Province; the upper contact is always disrupted by diapirism.
	*Coominaree Dolomite (Eok) Thomson, 1966 Coats and Blissett, 1971 Preiss, 1973	77	Interbedded buff dolomites (stromatolitic) and minor sandstones in the lower part; the upper part consists of non stromatolitic dolomites, oolitic at the base, and an overlying stromatolitic dolomite.	Contains stromatolite types <u>Acaciella c.f. australica</u> and <u>Gymnosolen c.f. ramsayi</u> .
	Younghusband Conglomerate (Eco) Thomson, 1966 Preiss, 1973	27	Basal quartzitic breccias, red brown shales and sandstones at the top.	Basal Adelaidean unit containing clasts reworked from the underlying Peake Metamorphics.
ANGULAR UNCONFORMITY				
PRE-ADELAIDEAN STRATIGRAPHY				
CARBONIFEROUS	*Wirriecurrie Granite (E8w) Reyner, 1955	-	Coarse-grained, porphyritic granites, augen granites, minor aplite dykes.	Intrusive, syn-orogenic granite, radiometrically dated at approximately 1600 Ma.
	Unnamed metamorphics (Ed3)	not known	Amphibolites, grey quartzites, magnetite-rich epidiosites, arenaceous schists, minor clinopyroxene granulites.	Lithologies (amphibolites and quartzites) are broadly similar to Baltucoodna Quartzite; all rock types are magnetite-rich.
PEAKE METAMORPHICS	Sedimentary contact not observed			
	*Baltucoodna Quartzite (Edb) Reyner, 1955	4500 (minimum)	Greyish-white quartzites, porphyritic and amygdaloidal basalts, minor quartz, muscovite schists, grey phyllites, sillimanite gneisses and plagioclase, hornblende, epidote calc-silicates.	Unit is characterised by interlayered basalts and quartzites.
	Unnamed schists (Ed2)	1000	Quartz, muscovite schists, grey quartz, chlorite phyllites, minor sandstones.	Thickest development of metamorphosed pelitic sediments.
	*Tidnamurkuna Volcanics (E8t) Preiss, 1973	600 (minimum)	Flow-banded porphyritic rhyolites, amygdaloidal basalts, rare epidiosites and phyllites.	Rhyolites record the only phase of acid volcanism in the Peake Metamorphics.
PEAKE METAMORPHICS	Sedimentary contact not observed			
	Unnamed metamorphics (Ed1)	5500 (minimum)	Grey quartzites, quartz, biotite and muscovite schists, gneisses, garnetiferous schists, migmatites, pegmatites and diorites.	Metamorphosed to amphibolite facies, sedimentary structures have been obliterated. Stratigraphic relationship with other units is not known.
TABLE 1. Summary stratigraphic table for the Peake and Denison Ranges, highlighting the lithologies, boundary relationships, depositional environments and unit thicknesses.				
* New names; for formal definition see Appendix A.				

APPENDIX A

Definition of new units

# APPENDIX A

## UNIT 1

NAME OF UNIT                      Tidnamurkuna Volcanics                      STATE    S.A.

PROPOSER                      R.B. Flint

### DEFINITION

DERIVATION OF NAME: Tidnamurkuna is an aboriginal name for a waterhole on the Peake Creek just west of the Peake and Denison Ranges, 9.5 km northwest of the ruins of the Peake overland telegraph station. Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area, metric reference 6898050, 580000.

DISTRIBUTION: The unit outcrops along a strike length of 10 km to the west of Peake ruins and 3 km north of Mount Denison on the WARRINA 1:250 000 sheet area.

TYPE SECTION: 500 metres of amygdaloidal basalts and porphyritic rhyolites exposed in the creek banks from 6888940, 590850 (bottom) to 6888300, 590750, (top of section). The base is not exposed and the top is identified by a cream weathering porphyritic rhyolite.

LITHOLOGY: Flow-banded porphyritic rhyolites, amgdaloidal basalts, minor epidotes and phyllites.

THICKNESS: Minimum thickness of 600 m.

RELATIONSHIPS AND BOUNDARY CRITERIA: The base is not exposed. Conformably overlain by unnamed schist unit of the Peake Metamorphics.

AGE AND EVIDENCE: ?Lower Proterozoic. The intrusive Wirriecurrie Granite has been radiometrically dated at  $1660 \pm 180 \text{ Ma}$  (Webb & Lowder, 1972; Webb, 1974) providing a minimum age.

## UNIT 2

NAME OF UNIT                      Baltucoodna Quartzite                      STATE    S.A.

PROPOSER                      R.B. Flint

### DEFINITION

DERIVATION OF NAME: Baltucoodna is an aboriginal name for a waterhole on the Peake Creek east of the Peake and Denison Ranges, 4.3 km north of the ruins of the Peake overland telegraph station. Warrina 1:100 000 sheet area, WARRINA 1:250 00 sheet area, metric reference: 6898300, 589350.

DISTRIBUTION: Peake and Denison Ranges; the unit is exposed over 25 km<sup>2</sup> south and southeast of Peake ruins and within various ? Lower Proterozoic basement rocks in the southern portion of the ranges.

TYPE SECTION: 4 500 metres of white quartzites and basalts from 6888100, 592200 (base) through Mount Denison 6886100, 594100 to 6883600, 592150 (top of section).

LITHOLOGY: In the type section, greyish-white quartzites, porphyritic and amygdaloidal basalts, minor quartz-muscovite schists and grey phyllites. Elsewhere also sillimanite gneisses, quartz-feldspar-biotite gneisses and schists, plagioclase-hornblende-epidote calc-silicates, calcite marble.

THICKNESS: Minimum of 4 500 metres.

RELATIONSHIPS AND BOUNDARY CRITERIA: In the type section the base is sheared, elsewhere the unit conformably overlies unnamed schists of the Peake Metamorphics. Top of the unit is not exposed.

AGE AND EVIDENCE: ?Lower Proterozoic. The intrusive Wirricurrie Granite has been radiometrically dated at  $1660 \pm 180$  Ma (Webb & Lowder, 1972, Webb, 1974) providing a minimum age.

### UNIT 3

NAME OF UNIT                      Bungadillina Monzonite                      STATE    S.A.

PROPOSER R.B.    Flint

### DEFINITION

DERIVATION OF NAME: Bungadillina is an aboriginal name for a waterhole and creek near Warrina railway siding. Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area, metric reference 6874650, 580950.

DISTRIBUTION: Occurs as sills, dykes, stocks and irregularly shaped intrusives around the headwaters of Bulldog, Levi and Bungadillina Creeks. Boorthanna and Anna Creek 1:100 000 sheet areas.

TYPE SECTION: Porphyritic monzonites, syenites and diorites from 6853100, 602250 to 6852700, 600000. Reference section of porphyritic monzonitic and syenitic sills from 6861400, 597050 to 6858500, 593400.

LITHOLOGY: Porphyritic monzonites and syenites, porphyritic diorites, porphyritic albitites, minor granites, adamellites, and granodiorites.

AGE AND EVIDENCE: Ordovician, radiometrically dated by K-Ar method (Webb, 1970; Webb and Lowder, 1971; 1972). Youngest unit intruded is the Torrensian Skillogalee Dolomite.

### UNIT 4

NAME OF UNIT    Younghusband Conglomerate                      STATE    S.A.

PROPOSERS G.J. Ambrose, R.P. Coats and W.V. Preiss

### DEFINITION

DERIVATION OF NAME: Mount Younghusband occurs within the Denison

Inlier in the northern Peake and Denison Ranges, 1.5 km east of Mount Denison Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area; metric ref: 6886500, 594500.

DISTRIBUTION: The unit outcrops along a strike length of 7 km, 3.5 km southwest of "Peake" ruins on Warrina. Other small outcrops occur 3 km south of War Loan Mine and 1.2 km north of Coominaree Mine.

TYPE SECTION: 27 metres of clastic sediments outcropping on the northeastern margin of the Coominaree Mine Block (?Lower Proterozoic basement), 1.2 km north of Coominaree Mine. The lower part (12 m thick) contains coarse clastic sediments. The upper 15 m is mainly argillaceous. Metric ref: Type Section 6853275, 596475. The unit was first described by Thomson (1966).

LITHOLOGY: Basal quartzitic breccia, red-brown shales and sandstones at the top.

THICKNESS: Variable up to 27 m (Preiss, 1973a).

RELATIONSHIPS AND BOUNDARY CRITERIA: Overlies ?Lower Proterozoic basement rocks (Peake Metamorphics) with angular unconformity. Reworked basement clasts recognised in the basal breccia. The unit is lenticular. In the type section it is conformably overlain by the Coominaree Dolomite while 3.5 km southwest of "Peake" ruins the Younghusband Conglomerate is disconformably overlain by the Cadlareena Volcanics.

AGE AND EVIDENCE: Basal Adelaidean - Callanna Beds (Willouran); an angular unconformity with Lower Proterozoic basement rocks is observed 3.5 km southwest of "Peake" ruins.

#### UNIT 5

NAME OF UNIT Coominaree Dolomite STATE S.A.

PROPOSERS G.J. Ambrose, R.P. Coats and W.V. Preiss

#### DEFINITION

DERIVATION OF NAME: Coominaree Mine situated 7 km northwest of Nilpinna H.S. in the Peake and Denison Ranges; Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area; metric reference is 6852300, 596080.

DISTRIBUTION: Small outcrops occur 3 km south of War Loan Mine and 1.2 km north of Coominaree Mine on the eastern margin of Margaret Inlier in the Peake and Denison Ranges.

TYPE SECTION: Occurs in a lower Adelaidean succession on the northeastern margin of the Coominaree Mine Block (?Lower Proterozoic); totals 77 metres in thickness; metric reference is 6853300, 596600. The unit was first described by Thomson (1966).

LITHOLOGY: The lower part of the unit consists of interbedded buff and pale brown dolomites with minor sandstones and pebbly layers; low angle cross-bedding is common. The upper part of the unit consists of a non stromatolitic dolomites, oolitic at the base, and an overlying stromatolitic unit.

THICKNESS: Variable up to 77 m.

RELATIONSHIPS AND BOUNDARY CRITERIA: A lenticular unit conformably overlying the Younghusband Conglomerate and in turn, conformably overlain by the Cadlareena Volcanics.

AGE AND EVIDENCE; Lower Adelaidean - Callanna Beds (Willouran); contains stromatolite types Acaciella c.f. australica and Gymnosolen c.f. ramsayi (Preiss, 1973a). In the Mount Painter Province the Wywyana Formation (Lower Callanna Beds) (Coats, 1971) is a carbonate-rich unit considered equivalent to the Coominaree Dolomite.

#### UNIT 6

NAME OF UNIT Cadlareena Volcanics

STATE S.A.

PROPOSERS G.J. Ambrose, R.B. Flint, R.P. Coats

#### DEFINITION

DERIVATION OF NAME: Cadlareena 1:63 360 sheet area, WARRINA 1:250 000 sheet area.

DISTRIBUTION: Peake and Denison Ranges; the unit is exposed over about 6 km<sup>2</sup> 1-2 km east of Douglas Well (metric ref: 6822020, 616750). A number of small outcrops occur on the eastern side of the Ranges about 5-10 km southeast of Nilpinna H.S. and 1.5 km northeast of Coominaree Mine. On Warrina about 3.5 km southwest of "Peake" ruins, the volcanics outcrop along a total strike length of 7 km.

TYPE SECTION: The basal part of the type section is on Warrina where a basal contact with the Younghusband Conglomerate is preserved, metric ref: base 6890725, 587562 - top 6890725, 687625. The upper part of the section occurs about 5 km east of Douglas Well on Anna Creek, metric ref: base 6823210, 622333 top 6823210, 622333.

THICKNESS: Maximum true thickness is uncertain. Maximum observed thicknesses are variable up to 750 m measured near Douglas Well.

LITHOLOGY: Vesicular basalts and altered dolerites; minor andesites, dacites and rhyolites; tuffs, lapilli tuffs; minor lenticular reddish mudstones and quartzites with red shale interbeds near base and top.

RELATIONSHIPS AND BOUNDARY CRITERIA: Disconformably overlies the Younghusband Conglomerate in the type section. Conformably overlies Coominaree Dolomite on the western side of the Peake and Denison Ranges. Throughout the Ranges diapirism occurs above the Volcanics and consequently the upper limit of the unit is unknown.

AGE AND EVIDENCE: Lower Adelaidean - Callanna Beds (Willouran); the Cadlareena Volcanics are the uppermost member of the Callanna Beds recognised in the Peake and Denison Ranges. In the Mount Painter Province the Wooltana Volcanics (Coats, 1971) are considered equivalent.



UNIT 7

NAME OF UNIT Rockwater Beds

STATE S.A.

PROPOSER G.J. Ambrose

## DEFINITION

DERIVATION OF NAME: Rockwater Hill situated 18 km southwest of Warrina R.S. on the Central Australia Railway; Umbum 1:100 000 sheet area. WARRINA 1:250 000 sheet area; metric ref: 6869200, 595510.

DISTRIBUTION: Several small outcrops occur as blocks in a diapir on the western margin of the Rockwater Hill Block. Two small blocks outcrop on Warrina about 11 km northeast of Warrina R.S. A third block outcrops immediately south of Coominaree Mine on Warrina.

TYPE SECTION: The unit only outcrops in blocks stratigraphically isolated by faulting and diapirism and hence a type section has not been defined. The best reference section is a block immediately south of Coominaree Mine on Warrina. Metric ref: 6852050, 595900.

THICKNESS: True maximum thickness is uncertain. The thickest section ( 200 m) is in the reference section south of Coominaree Mine; metric ref: 6852050, 595900.

LITHOLOGY: Blue-grey and black cherts, black dolomitic limestones, black silty shales and grey quartzites.

RELATIONSHIPS AND BOUNDARY CRITERIA: The unit is stratigraphically isolated by faulting and diapirism; thought to occur stratigraphically above the Cadlareena Volcanics and below the War Loan Beds although no contacts are observed.

AGE AND EVIDENCE: Lower Adelaidean - it is not certain whether the unit is Willouran or Torrensian; the unit probably closely overlies the Willouran Cadlareena Volcanics although no contact is exposed; it correlates with the Dunn's Mine Beds of the Willouran Ranges (Murrell, 1977) which closely overly Willouran volcanics in that region.

UNIT 8

NAME OF UNIT War Loan Beds

STATE S.A.

PROPOSERS G.J. Ambrose and R.P. Coats

## DEFINITION

DERIVATION OF NAME: War Loan Mine situated on the western margin of the Peake and Denison Ranges, 8.5 km southeast of Nilpinna H.S., Boorthanna 1:100 000 sheet area, WARRINA 1:250 000 sheet area; metric ref: 6844733, 597225.

DISTRIBUTION: Outcrops in a synclinal structure adjacent to War Loan Mine.

TYPE SECTION: 600 m of sediments outcropping in a synclinal structure adjacent to War Loan Mine; metric reference - base 6844675, 597225 - top 6844210, 597233.

LITHOLOGY: Blue-grey shales and siltstones, dark grey silty dolomites, greenish feldspathic sandstones and arkoses.

RELATIONSHIPS AND BOUNDARY CRITERIA: Faulted against the Cadlarena Volcanics in its type section. Probably grades upwards into the Nilpinna Beds although all contacts are tectonic.

THICKNESS: Maximum true thickness uncertain; the thickest development of sediments, 600 metres, is in the type section.

AGE AND EVIDENCE: Lower Adelaidean (Willouran or Torrensian). The unit is believed to grade upwards into the Nilpinna Beds; lithological affinities and spatial relationships between outcrops of these units support this view. The lower contact is uncertain but probably occurs somewhere above the Rockwater Beds.

#### UNIT 9

NAME OF UNIT Nilpinna Beds

STATE S.A.

PROPOSERS G.J. Ambrose and R.P. Coats

#### DEFINITION

DERIVATION OF NAME: Nilpinna H.S. situated on the eastern margin of the Peake and Denison Ranges, 6.5 km east of Duff Creek R.S. on the Central Australia Railway; Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet Area. Metric reference is 6849133, 5900075.

DISTRIBUTION: Outcrops in a west-facing, fault-bound sequence 1 km north of War Loan Mine; metric ref: 6844733, 597225.

TYPE SECTION: 2 100 metres of sediments in a fault bound sequence outcropping 6 km southeast of Nilpinna H.S. The base of the section (metric ref: 6846125, 596650) is faulted against War Loan Beds and the top of the section (metric ref: 6846175, 594650) is faulted against Duff Creek Beds.

LITHOLOGY: Fine cross-laminated, rippled sandstones, rippled quartzites (salt casts), green-grey silty shales, minor grey silty dolomites and greenish arkoses; mudcracks, ripple marks and salt casts are common.

THICKNESS; Maximum true thickness is uncertain. The thickest development of sediments (2100 m) is in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: Believed to grade downwards into the War Loan Beds and upwards into the Duff Creek Beds although all contacts with these units are sheared. Lithological affinities and spatial distribution of outcrops support this view.

AGE AND EVIDENCE: Lower Adelaidean (Willouran or Torrensian); probably intervenes between the War Loan Beds and Duff Creek Beds.

UNIT 10

NAME OF UNIT Duff Creek Beds

STATE S.A.

PROPOSERS G.J. Ambrose (after R.P. Coats and B.P. Thomson, 1964)

## DEFINITION

DERIVATION OF NAME: Duff Creek R.S. on the Central Australia Railway, Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area. Metric reference is 68498500, 583730.

DISTRIBUTION: Outcrops in several thick sequences on the eastern margin of the Peake and Denison Ranges, east and south of Nilpinna H.S., and in faulted sequences 7 km southeast of "Peake" ruins and immediately east of Mount Kingston. Other faulted blocks containing Duff Creek Beds occur west of Tarlton Springs, 5.5 km northwest of Douglas Well, 4 km west of Mount Denison and one kilometre south of Mount Dutton.

TYPE SECTION: The type section is defined by two separate sections totalling 5 500 m in thickness. The upper part of the type section (3 700 m) occurs east of Nilpinna H.S. and is defined by metric refs: base 6847600, 595050 - top 6849025, 5911660, The lower 1 800 m of the type sequence outcrops 9 km to the south and is defined by metric refs: base 6838350, 593700 - top 6841090, 592960.

LITHOLOGY: Laminated olive green silty shales and thin arkose interbeds, flaggy buff-weathering pale grey and yellow dolomites, minor dark grey dolomites and carbonaceous shales, flaggy dolomitic siltstones, laminated pyritic fine sandstones, minor quartzitic sandstones; mudcracks, ripple marks, clay galls and cross-bedding are common; horizons with abundant salt casts and gypsum casts; algal laminations.

THICKNESS: Maximum true thickness is uncertain. Thickest development of sediments (5 500 m) is in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: Exhibits faulted contacts with underlying and overlying sequences. Believed to grade downwards into the Nilpinna Beds and upwards into the Murrana Beds although all contacts with these units are sheared. Lithological affinities and spatial distribution of outcrops support this view.

AGE AND EVIDENCE: Lower Adelaidean (Willouran or Torrensian). Probably intervenes between the Nilpinna Beds and Murrana Beds NB. Reyner (1955) defined the Duff Creek Formation to include all Willouran rocks in the Peake and Denison Ranges.

UNIT 11

NAME OF UNIT Murrana Beds

STATE S.A.

PROPOSER G.J. Ambrose

## DEFINITION

DERIVATION OF NAME: Murra Murrana bore situated on the southern

margin of the Denison Inlier in the Peake and Denison Ranges, 5 km northeast of Peake H.S. (metric ref: 6878480, 593250); Warrina 1:100 000 sheet area; WARRINA 1:250 000 sheet area.

DISTRIBUTION: Outcrops in a basinal syncline 4 km northwest of Warrina R.S.; constitutes most of Mount Dutton Inlier; outcrops in a faulted sequence 1 km east of Nilpinna H.S.

TYPE SECTION: Comprises 2 900 m of sediments outcropping on the eastern limb of a basinal syncline 4 km northwest of Warrina R.S. Metric reference of the base of the section is 6885100, 587650; of the top 6885680, 581290. Contacts at the base and top of the section are faulted.

LITHOLOGY: Laminated gritty quartzitic sandstones, grey-green silty shale interbeds; arkoses, pebbly dolomites, purple and grey silty shales near the top; ripple marks, mud cracks, cross-laminations flute casts and salt casts.

THICKNESS: Maximum true thickness is uncertain. Thickest development of sediments (2 900 m) is in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: Nature of overlying sequences is uncertain. The unit is often faulted at the base against Duff Creek Beds. Lithological affinities with the Duff Creek Beds suggest a gradational contact between the two units although this has not been observed intact in the field.

AGE AND EVIDENCE: Lower Adelaidean (Willouran or Torrensian). The unit probably grades down into the Duff Creek Beds.

#### UNIT 12

NAME OF UNIT Fountain Spring Beds

STATE S.A.

PROPOSER G.J. Ambrose

#### DEFINITION

DERIVATION OF NAME: Fountain Spring on the plains abutting the eastern margin of the Peake and Denison Ranges, 43 km east of Edwards Creek R.S. on the Central Australia Railway. Umbum 1:100 000 sheet area, WARRINA 1:250 000 sheet area. Metric ref: 6863290, 625690.

DISTRIBUTION: Outcrops solely on the middle-eastern and northwestern margins of the Margaret Inlier in the Peake and Denison Ranges.

TYPE SECTION: 1 100 metres of sediments occur in the type section 3 km north of Mount Margaret, metric reference: base 6853033, 603590 - top 6852020, 603590.

LITHOLOGY: Laminated grey dolomitic siltstones and interbedded thick grey quartzites with clay gall laminations, grey-green silty shales (salt casts); minor grey silty dolomites and a few quartzites near the base.

THICKNESS: Maximum true thickness is uncertain. Thickest development of sediments (1 100 m) is in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: Grades upwards into the Mount Margaret Quartzite the contact being marked by an increase in orthoquartzites. The lower contact is tectonic although the unit probably grades downwards into an unnamed siltstone with which it has many lithological affinities.

AGE AND EVIDENCE: Adelaidean - Burra Group (Torrensian); considered equivalent to the River Wakefield Subgroup (Forbes, 1964).

### UNIT 13

NAME OF UNIT Mount Margaret Quartzite STATE S.A.

PROPOSERS G.J. Ambrose (after Reyner (1955) and Thomson and Coats (1964)).

### DEFINITION

DERIVATION OF NAME: Mount Margaret, on the eastern margin of the Margaret Inlier in the Peake and Denison Ranges, 15.5 km east of Nilpinna H.S., Umbum 1:100 000 sheet area, WARRINA 1:250 000 sheet area - metric reference 6848490, 605010.

DISTRIBUTION: The formation is restricted to the southern part of the Peake and Denison Ranges (i.e. the Margaret Inlier) and outcrops at Mount Margaret and also constitutes the Mount Margaret Plateau.

TYPE SECTION: Outcrops 5.5 km north-northwest of Mount Fox. From the base of the section (metric ref: 6853810, 598120) to the top (metric ref: 6850175, 600675) measures 2500 m.

LITHOLOGY: White orthoquartzites, slaty quartzitic sandstones, dark grey sandy siltstones, green-grey silty shales, minor dolomitic siltstones near the base; thick orthoquartzites at the top; clay galls, ripple marks and cross-bedding.

THICKNESS: Maximum total thickness of the formation is 2 500 m in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: The lower contact with the Fountain Springs Beds is gradational. A high silt-shale: quartzite ratio marks the transition down into the Fountain Spring beds. The Mount Margaret Quartzite is conformably overlain by a sandy facies of the Skillogalee Dolomite.

AGE AND EVIDENCE: Adelaidean - Burra Group (Torrensian).

SYNONYMS: Mount Margaret Quartzites (Reyner, 1955), Mount Margaret Formation (Thomson and Coats, 1964); Mount Margaret Quartzite (Thomson in Parkin, 1969).

UNIT 14

NAME OF UNIT Kalachalpa Formation

STATE S.A.

PROPOSER G. Ambrose

## DEFINITION

DERIVATION OF NAME: Kalachalpa is the Aboriginal synonym for Anna Creek which is a creek on the southern margin of the Peake and Denison Ranges. Anna Creek also gives its name to a cattle station encompassing most of the Ranges and a railway siding on the Central Australia Railway; Boorthanna 1:100 000 sheet area; WARRINA 1:250 000 sheet area.

DISTRIBUTION: Outcrops in a southwesterly facing sequence, 6 km north of Box Creek R.S. on the Central Australia Railway. A faulted sequence of the formation outcrops 1 km northeast of Box Creek R.S. Smaller outcrops occur 8 km northwest and 11 km north of Anna Creek H.S.

TYPE SECTION: The type section occurs in an area first described in detail by Fairchild (1975). The type section outcrops 6 km north of Box Creek R.S., metric ref. base 6826575, 595450 - top 6824900, 594275; thickness is 900 metres.

LITHOLOGY: Grey-green and brown siltstones and shales, gritty sandstones and quartzites, conglomeratic dolomites, stromatolitic dolomites, oolitic sediments, black cherts; quartzites and shales at the top; abundant sedimentary features throughout the sequence include ripple marks, cross-bedding, graded bedding, clay galls and mud cracks.

THICKNESS: Maximum total thickness of the formation is 900 metres in the type section.

RELATIONSHIPS AND BOUNDARY CRITERIA: Grades downwards into an unnamed transitional unit which in turn overlies the Skillogee Dolomite. The formation is disconformably overlain by Calthorinna Tillite.

AGE AND EVIDENCE: Adelaidean-Uppermost Burra Group (Torrensian); probably correlates with the Myrtle Springs Formation (Coats, 1973).

UNIT 15

NAME OF UNIT Calthorinna Tillite

STATE S.A.

PROPOSERS G. Ambrose and R.P. Coats

## DEFINITION

DERIVATION OF NAME: Calthorinna Creek which is a tributary of Anna Creek and joins the main stream 1 km south of Anna Creek R.S. on the Central Australia Railway; Boorthanna 1:100 000 sheet area; WARRINA 1:250 000 sheet area; metric ref: 6812550, 607475.

**DISTRIBUTION:** The unit is restricted in outcrop to a syncline 1 km northeast of Box Creek R.S. (Box Creek Syncline) in the southern part of the Peake and Denison Ranges (Margaret Inlier). A smaller outcrop occurs as a rafted block in a diapir 6 km south of Mount Anna.

**TYPE SECTION:** 650 metres of sediments outcropping on the northern limb of the Box Creek Syncline, 5.5 km northwest of Box Creek R.S. The top of the unit is not exposed. Metric ref: Base 6825850, 591950 - top 6824710, 591550.

**LITHOLOGY:** Diamictites, conglomeratic dolomites, laminated green shales, gritty sandstones, arkoses and quartzites; cross-bedding, ripple marks and graded bedding are common. The unit was first described by Reyner (1955).

**THICKNESS:** Maximum thickness of the unit observed (650 m) is in the type section. True maximum thickness of the unit is uncertain since the top of the unit is not exposed in the type section.

**RELATIONSHIPS AND BOUNDARY CRITERIA:** The unit is underlain disconformably by upper Burra Group sediments (Kalachalpa Formation). It is overlain by and interfingers with an unnamed sandstone.

**AGE AND EVIDENCE:** Adelaidean - Umberatana Group (Sturtian). Believed to represent the second of the two Sturtian glaciations; main evidence is the presence of gritty quartzite clasts assumed to have been reworked from the Pandurra Formation (Coats and Forbes, 1977).

#### UNIT 16

NAME OF UNIT Thora Dolomite

STATE S.A.

PROPOSERS G. Ambrose and R.P. Coats

#### DEFINITION

**DERIVATION OF NAME:** Thora Soakage, 2 km northeast of Peake H.S. in the Peake and Denison Ranges; Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area; metric reference 6877620, 589800.

**DISTRIBUTION:** Sole outcrop is on the southern limb of a syncline (Box Creek Syncline) 5 km northeast of Box Creek R.S. on the Central Australia Railway; Boorthanna 1:100 000 sheet area, WARRINA 1:250 000 sheet area.

**TYPE SECTION:** 30-40 m of sediments on the southern limb of the Box Creek Syncline 5 km northeast of Box Creek R.S., metric reference 6820525, 601350; section occurs where a tributary of Wilyalallina Creek cuts through the units.

**LITHOLOGY:** Grey-green and brown dolomites, minor siltstones; weathers a light brown buff colour; large festoon surfaces and algal bedding.

THICKNESS: Range 30-40 metres.

RELATIONSHIPS AND BOUNDARY CRITERIA: Conformably overlies the Tapley Hill Formation (Coats, 1971); gradational upper contact with an unnamed siltstone of the Willochra Subgroup.

AGE AND EVIDENCE: Adelaidean Umberatana Group (Sturtian).

#### UNIT 17

NAME OF UNIT Wirriecurrie Granite STATE S.A.

PROPOSER R.B. Flint

#### DEFINITION

DERIVATION OF NAME: Wirriecurrie is the aboriginal word for Peake Creek which cuts the ranges north of the ruins of the Peake overland telegraph station. Warrina 1:100 000 sheet area, WARRINA 1:250 000 sheet area.




DISTRIBUTION: The granite outcrops in an area of 15 km<sup>2</sup> to the west and northwest of Peake ruins and 4 kms north of Mount Denison on the Warrina 1:250 000 sheet area.

TYPE AREA: 3 km northwest of Peake ruins, area bounded by metric references 6896600, 6894700 and 585800, 587800.

LITHOLOGY: Coarse-grained porphyritic granites, augen granites, adamellites, granodiorites, minor aplitic dykes.

AGE AND EVIDENCE: Middle Proterozoic, radiometrically dated by K-Ar method on biotite yielding an age of 1050 Ma and by Rb-Sr method on total rock giving an age of 1660 ± 180 Ma (Webb and Lowder, 1972; Webb, 1974). Intrudes the Tidnamurkuna Volcanics and unnamed metamorphics north of Peake Creek.







APPENDIX B

DEPARTMENT OF MINES AND ENERGY SOUTH AUSTRALIA		SCALE:
COMPILED: G. Ambrose		DATE: 8/3/78
DRN: P.D	CKD:	PLAN NUMBER
PEAKE AND DENISON RANGES STRATIGRAPHIC TYPE SECTIONS CALLANNA BEDS		78 - 211


## SECTION 4




## SECTION 5




## SECTION 6B




## SECTION 6A



## SECTION 6B ctd




## SECTION 7




## APPENDIX C

DEPARTMENT OF MINES AND ENERGY SOUTH AUSTRALIA		SCALE
COMPILED G. Ambrose		DATE 8/ 3/ 78
DRN P. D.	CKD	PLAN NUMBER
		78-214


SECTION 8



SECTION 9 (after Cooper et al, 1963)




SECTION 10 (after Cooper et al, 1963; Fairchild, 1975)




APPENDIX D

DEPARTMENT OF MINES AND ENERGY — SOUTH AUSTRALIA			Scale
Compiled: G.Ambrose			Date: 17/2/78
Drn. P.D.	Ckd.	PEAKE AND DENISON RANGES STRATIGRAPHIC TYPE SECTIONS BURRA GROUP	
			Org. No. 78-213

SECTION 11 Northern Limb of the Box Creek Syncline



SECTION 12 Southern Limb of the Box Creek Syncline



APPENDIX E

DEPARTMENT OF MINES AND ENERGY SOUTH AUSTRALIA		SCALE
COMPILED G. Ambrose		DATE 8/3/78
DRN P.D	CKD	PLAN NUMBER
PEAKE AND DENISON RANGES STRATIGRAPHIC TYPE SECTIONS UMBERATANA GROUP		78-212

- APPENDIX B - Stratigraphic type sections for new units  
in the Callanna Beds.
- APPENDIX C - Stratigraphic type sections for new units  
in Sequences of uncertain age.
- APPENDIX D - Stratigraphic type sections for new units  
in Burra Group.
- APPENDIX E - Stratigraphic type sections for new units  
in the Umberatana Group.







FIG.1

COMPILED, R.B. Flint  
 CHN. L.P.R. AND, R.H.

DEPARTMENT OF MINES AND ENERGY  
 SOUTH AUSTRALIA  
 PEAKE AND DENISON RANGES  
 LOCALITY PLAN

DATE August 1977  
 PLAN NUMBER  
 S13007

JOB NO. 44



S.A. DEPT. OF MINES AND ENERGY

PEAKE and DENISON RANGES - COOBER PEDY REGION

LANDSAT IMAGE 1208-00214-5

SCALE 1:1000 000

FIG. 2

S 13265








FIG. 4. Vesicle and infilling quartz and fibrous epidote in amygdaloidal basalt (Tidnamurkuna Volcanics)  
Photo No. 30296





FIG. 5. Plagioclase and quartz phenocrysts within porphyritic rhyolite (Tidnamurkuna Volcanics)  
Photo no. 30297




FIG. 6. Phenocrysts of feldspars and quartz within porphyritic rhyolite (Tidnamurkuna Volcanics)  
The pen is 150 mm long  
Photo no. 30298




FIG. 7.  
Plagioclase phenocrysts  
in basalts within the  
Baltucoodna Quartzite.  
Photo No. 30299




FIG. 8  
Heavy mineral banding  
in quartzites near  
Spring Hill.  
Photo No. 30300



FIG. 9  
Plagioclase and potash  
feldspar augen in the  
Wirriecurrie Granite.  
Photo No. 30301



FIG. 10 Younghusband  
Conglomerate capping  
Peake Metamorphics  
11.5 km northnortheast of  
Warrina R.S.  
Photo No. 30302




FIG. 11. Angular  
unconformity between  
steeply dipping meta-  
siltstones of the Peake  
Metamorphics and over-  
lying, gently dipping  
Cadlareena Volcanics,  
11 km northeast of  
Warrina R.S.  
Photo No. 30303



FIG. 12. Interbedded  
vesicular basalt and  
lenticular mudstone in  
the Cadlareena Volcanics.  
Photo No. 30304






FIG. 14. Well developed cross-laminations in the lower part of the Nilpinna Beds, 1.5 km north-west of War Loan Mine.

Photo No. 30305



FIG. 15. Typical laminated, buff weathering dolomite in the Duff Creek Beds, 6 km northwest of Douglas Well; note the thin siliceous interbeds.

Photo No. 30306



FIG. 16. Mud cracks in a silt dolomite of the Duff Creek Beds, west of Tarlton Springs.

Photo No. 30307



FIG. 17. Rugged, faulted, eastern margin of Margaret Inlier, north of Tarlton Springs. Quartzites of the Mount Margaret Quartzite are faulted against Mesozoic and younger sediments. Mount Margaret is situated right of centre on the skyline. Photograph by G.M. Pitt  
Photo No. 30308



FIG. 18. Looking south on the western margin of Margaret Inlier. Quartzites of the Mount Margaret Quartzite are faulted on the right against (topographically lower) siltstones of the Fountain Spring Beds. The penneplain surface forming the Mount Margaret Plateau dips gently to the west. Photograph by L.C. Barnes.  
Photo No. 30309




FIG. 19. Looking northwest-wards across gently dipping dolomites and sandstones of the Skillogalee Dolomite in the Margaret Inlier. The Mount Margaret Plateau is visible on the skyline. Photograph by G.M. Pitt.  
Photo No. 30310



Fig. 21. Intraformational chert pebble-cobble conglomerate in the upper part of the unnamed transition unit, Boorthanna Section. Photo No. 30311



Fig. 22. Conglomeratic dolomite typical of much of the upper Burra Group, showing convolute bedding and graded bedding. Most of the clasts are intraformational. Photo No. 30312




Fig. 23. Columnar stromatolites in the Kalachalpa Formation on the southern limb of the Box Creek Syncline. Photo No. 30313





FIG. 24. Conglomeratic dolomite overlying a more massive dolomite in the Calthorinna Tillite on the southern limb of the Box Creek Syncline. The conglomerate contains dolomite clasts reworked from the underlying dolomite. Photo No. 30314



FIG. 25. Sharp contact between a conglomeratic dolomite (left) and diamictite (right) in the Calthorinna Tillite on the southern limb of the Box Creek Syncline. The diamictite has a quartz rich matrix. Photo No. 30315




FIG. 26. Reworked dark coloured diamictite clasts set in a pale dolomite, in the upper part of the Calthorinna Tillite type section on the northern limb of the Box Creek Syncline. Chert and dolomite clasts are also common in this unit. Photo No. 30316



FIG. 27  
Feldspar phenocrysts in  
porphyritic monzonite of  
the Bungadillina Monzonite.  
The pen is 150 mm long.  
Photo No. 30317



FIG. 28  
Aligned feldspar phenocrysts  
in a monzonite of the  
Bungadillina Monzonite.  
Photo No. 30318




FIG. 29  
Zoned plagioclase phenocryst  
in a syenite of the Bungadi-  
llina Monzonite.  
Photo No. 30319









FIG. 30. Angular unconformity between dipping dolomites of the Kalachalpa Formation and overlying, flat lying Permian boulder lag, 6 km north of Box Creek R.S. The boulder lag (Boorthanna Formation) contains boulders of mainly local Adelaidean sediments. Photo No. 30320



FIG. 31. Typical view of brecciated and highly contorted lower Adelaidean sediments in a broad diapiric zone above the Cadlareena Volcanics 9 km northeast of Warrina R.S. The carbonate matrix breccia between the blocks outcrops poorly. Photo No. 30321



FIG. 32. Diapiric, carbonate matrix breccia; deformed blocks of Adelaidean sediments in a carbonate matrix. Photo No. 30322



# GEOLOGICAL MAP PEAKE AND DENISON RANGES

SCALE 1:100 000




## REFERENCE

### TOPOGRAPHICAL SYMBOLS


- Road
- Main
- Secondary
- Track
- Railway
- with station
- with siding
- Triangulation station
- Identified point
- Homestead, building
- Yard
- Mine
- Effluential stream
- Flood plain boundary
- Swamp
- Spring
- Waterhole
- Dam
- Bore, well, tank
- Sand dunes


### GEOLOGICAL SYMBOLS

- Geological boundary
- Observed
- Inferred
- Fault
- Observed
- Inferred
- Fold
- Anticline
- Syncline
- Bedding
- Inclined
- Vertical
- Trend of bedding
- Folding
- Foliation
- Vertical
- Geological block diagram
- Type section



Geology by G.J. Ambrose and R.B. Flint






**PLAN No. 78-868**








S.M.L. 84  
S.M.L. 125 Portion of S.M.L. 255 above 28° 27' (28° 27' shown thus — — — — —)  
S.M.L. 134 Portion of S.M.L. 255 below 28° 27'  
S.M.L. 255  
S.M.L. 270  
S.M.L. 329  
S.M.L. 437  
S.M.L. 492  
S.M.L. 546 Same boundary as S.M.L. 270  
S.M.L. 718

Area of inliers


TENEMENT NUMBER	S.A. DIM ENV. No.	HOLDER	ELEMENTS		STRATIGRAPHY	MAJOR EXPLORATION METHODS	TENURE	
			MAJOR	MINOR			COMMENCED	EXPIRED
S.M.L. 84	No. record available	K. Wright					1-7-65	6-10-66
S.M.L. 125	941	North Broken Hill Ltd	Cu	Pb, Zn, Ni	Precambrian	C P D	1-10-66	30-9-68
S.M.L. 134	941	North Broken Hill Ltd	Cu	Pb, Zn, Ni	Precambrian	C P D	1-11-66	31-10-68
S.M.L. 255	941	North Broken Hill Ltd	Cu	Pb, Zn, Ni	Precambrian	C P D	1-11-66	30-10-69
S.M.L. 270	1015	Australasian Mining Corp. Ltd.	Cu		Precambrian	P G D	17-2-69	16-2-71
S.M.L. 329	1241, 1313	Occidental Minerals Corp.	K	S, coal	Palaeozoic	P D	14-8-69	13-8-71
S.M.L. 437	1471	Stockholm Mining N.L.	Clay		Mesozoic	G	18-6-70	18-6-71
S.M.L. 492	1015	Australasian Mining Corp. Ltd.	Cu		Precambrian	P G D	5-11-70	4-5-71
S.M.L. 546	1015	Australasian Mining Corp. Ltd.	Cu		Precambrian	G	25-2-71	24-2-72
P.E.L. 5/6	1374, 1495	Pexa Oil N.L.	Hydrocarbons		Palaeozoic	P D H	28-2-69	21-6-73
E.L. 22	2182	Chevron Exploration Corp.	U		Mesozoic	G D	2-11-72	1-11-73
E.L. 33	2253	Uranerz (Aust.) Pty. Ltd.	U	Cu	Precambrian	G	4-1-73	3-1-74
E.L. 108	2388	Shell Development (Aust.) Pty. Ltd.	Coal		Palaeozoic	P D H	1-11-73	31-10-74
E.L. 110	2381	Uranerz (Aust.) Pty. Ltd.	U		Precambrian	C P D	16-11-73	15-11-74
E.L. 192	2525	Western Mining Corp.	Au, Cu	Ni, Co, Cr, Ag, Mn, Pb, Zn, As	Precambrian	C G	5-5-75	4-5-76
E.L. 336	confidential	Mines Administration Pty. Ltd. & Teton Exploration Drilling Co.					12-7-77	
E.L. 340	confidential	Newmont Pty. Ltd. & Dampier Mining Co. Ltd.					10-8-77	
E.L. 369	confidential	Dampier Mining Co. Ltd.					31-10-77	
S.M.L. 718	2142	A.O.G. Minerals Pty. Ltd.	Au		Mesozoic	G	22-6-72	21-6-73



E.L. 22  
E.L. 33  
E.L. 108  
E.L. 110  
E.L. 192  
E.L. 336  
E.L. 340  
E.L. 369

#### LEGEND

S.M.L. Special Mining Lease  
P.E.L. Petroleum Exploration Licence  
E.L. Exploration Licence  
C Geochemistry  
P Geophysics  
G General  
D Drilling  
H Hydrocarbons



#### Enclosure 3

DEPARTMENT OF MINES AND ENERGY - SOUTH AUSTRALIA

### PEAKE AND DENISON RANGES MINING TENEMENTS

COMPILED G. J. A.	DRN I.M.	SCALE 1:500 000	PLAN NUMBER
DIRECTOR GENERAL	CKE	DATE 3-4-78	78-316

ROCK SPECIMEN NUMBER	ANALYSES NUMBER	ROCK NAME	LITHOLOGICAL UNIT	ELEMENTS																							
				Ag	As	Au	Ba	Bi	Co	Cr	Cu	Li	Mn	Mo	Ni	P	Pb	Rb	Sb	Sn	Sr	Th	U	V	W	Zn	Zr
6040 RS 14	A 3378/63	DOLERITE	CADLAREENA VOLCANICS	0.3					50	700	40		400		120		12							600		120	
6040 RS 15	A 3379/63	"	"	0.2						1000	10		600		120		15							500		180	
6040 RS 19	A 4860/76	GNEISS	BALTUCOODNA QUARTZITE	X	X	X	1500	X	100	250	150		500	10	200	500	3		X	X				300	X	50	
6040 RS 20	A 4861/76	GRANULITE	"	X	X	X	X	X	20	80	10		70	X	70	1000	15		X	1				50	X	X	
6040 RS 21	A 4862/76	GNEISS	"	X	X	X	1000	X	80	150	30		1000	10	100	500	3		X	5				300	X	X	
6041 RS 12	A 4879/76	GRANITIC GNEISS	WIRRIE CURRIE GRANITE			+	1200		45	20	30	30		+	10		75	250				140	22	12	30	45	340
6041 RS 43	A52/72	DOLERITE	DYKE;						80	100	40																
6041 RS 44	A53/72	MONZONITE	BUNGADILLINA MONZONITE						50		20			X						1					X		
6041 RS 45	A54/72	"	"						30		10			X						1					X		
6041 RS 46	A55/72	"	"						30		10			X						1					X		
6041 RS 58	A 4878/76	SYENITE	"			+	945		45	20	10	10		+	10		20	120				1050	6	N.D.	80	15	135
6041 RS 62	A 4880/76	GRANITE PORPHYRY	WIRRIE CURRIE GRANITE			+	1250		55	15	95	35		10	10		28	250				250	36	8	35	60	450
6041 RS 93	A30/76	DACITE	CADLAREENA VOLCANICS	0.5				X	X	30	80					5		800					N.D.	N.D.	100	X	
6041 RS 95	A31/76	DOLERITE	"	X				X	100	300	10					150		20					N.D.	N.D.	500	30	
6041 RS 96	A32/76	"	"	X				X	50	200	20					80		10					N.D.	N.D.	500	X	
6041 RS 97	A33/76	BASALTIC PYROCLASTIC	"	X				X	10	100	5					30		10					N.D.	8	300	X	
6041 RS 98	A34/76	ANDESITE	"	X				5	X	80	10					30		10					N.D.	22	200	X	
6041 RS 99	A35/76	DOLERITE	"	X				X	X	20	1				X		10						N.D.	N.D.	10	50	
6041 RS 100	A36/76	ANDESITE	"	X				X	30	100	5					20		10					4	30	200	X	
6041 RS 101	A37/76	BASALT	"	X				X	50	80	5					20		10					N.D.	N.D.	200	20	
6041 RS 114	A785/76	MONZONITE	BUNGADILLINA MONZONITE	X		+			5	20	50			X	10		3			X			12	4	150	X	
6041 RS 115	A786/76	SYENITE	"	X		+			5	30	150			X	10		5			X			16	N.D.	100	X	
6041 RS 116	A797/76	MONZONITE	"	X		+			5	20	10			X	10		3			X			16	N.D.	80	X	
6041 RS 124	A4845/76	CALC-SILICATE	BALTUCOODNA QUARTZITE	X	X	X	800	X	80	100	50		500	X	150	500	10		X	1				150	X	50	
6041 RS 125	A4847/76	GNEISS	UNNAMED METAMORPHICS Ed <sub>1</sub>	X	X	X	700	X	30	70	30		300	X	10	1000	3		X	X				150	X	X	
6041 RS 126	A4848/76	SCHIST	"	X	X	X	500	X	100	100	150		800	X	200	300	1		X	X				200	X	50	
6041 RS 127	A788/76	MICROGRANITE	BUNGADILLINA MONZONITE	X		+			5	20	5			X	5		5			X			16	6	80	X	
6041 RS 128	A794/76	DIORITE	"	X		+			X	20	10			X	X		5			X			14	N.D.	10	X	
6041 RS 129	A4849/76	CALC-SILICATE	BALTUCOODNA QUARTZITE	X	X	X	X	X	80	100	30		1000	10	80	300	10		X	X				200	X	150	
6041 RS 130	A4850/76	SCHIST	"	X	X	X	1000	1	80	150	30		300	X	150	500	3		X	2				200	X	50	
6041 RS 135	A4851/76	METABASALT	TIDNAMURKUNA VOLCANICS	X	X	X	300	X	200	300	100		1000	X	300	500	3		X	X				500	X	X	
6041 RS 136	A4852/76	"	"	X	X	X	1000	X	150	300	100		800	X	250	1000	10		X	X				300	X	100	
6041 RS 137	A4853/76	AMPHIBOLITE	BALTUCOODNA QUARTZITE	X	X	X	X	X	30	100	200		1000	X	80	200	10		X	X				100	X	150	
6041 RS 138	A4854/76	METABASALT	TIDNAMURKUNA VOLCANICS	X	X	X	700	X	200	200	300		1000	X	300	500	3		X	X				300	X	70	
6041 RS 139	A4855/76	"	BALTUCOODNA QUARTZITE	X	X	X	500	X	150	200	500		1000	X	300	1000	5		X	X				500	X	X	
6041 RS 140	A4856/76	AMPHIBOLITE	DYKE;	X	X	X	500	X	80	150	70		1000	X	100	1000	10		X	X				200	X	80	
6041 RS 141	A4857/76	"	WIRRIE CURRIE GRANITE	X	X	X	700	X	200	150	100		800	X	200	500	5		X	X				500	X	50	
6041 RS 142	A4859/76	DIORITE	BALTUCOODNA QUARTZITE	X	X	X	1500	X	200	300	80		500	X	250	X	3		X	X				300	X	50	
6041 RS 143	A4876/76	METADOLERITE	WIRRIE CURRIE GRANITE			+	300		55	120	170	15		+	100		10	50				95	N.D.	14	170	90	165
6041 RS 144	A4873/76	RHYOLITE	BALTUCOODNA QUARTZITE			+	960		70	10	45	15		+	15		32	210				44	34	8	+	25	570
6041 RS 145	A4874/76	"	TIDNAMURKUNA VOLCANICS			+	1000		80	15	15	10		+	10		26	260				65	30	12	20	20	240
6041 RS 146	A4875/76	DACITE	"			+	800		55	90	240	25		+	100		10	95				100	8	N.D.	210	45	165
6041 RS 147	A4863/76	AMPHIBOLITE	UNNAMED METAMORPHICS Ed <sub>1</sub>	X	X	X	500	X	30	100	30		100	X	100	5000	1		X	1					X	X	
6041 RS 148	A4864/76	"	"	X	X	X	300	X	150	800	200		1000	X	300	1000	10		X	1					X	80	
6041 RS 149	A4865/76	"	"	X	X	X	700	X	200	100	200		1000	X	200	500	5		X	1					X	100	
6041 RS 150	A4866/76	SCHIST	UNNAMED SCHISTS Ed <sub>2</sub>	X	X	X	800	X	150	300	100		500	20	300	500	3		X	X				200	X	X	
6041 RS 151	A4867/76	"	"	X	X	X	200	X	100	100	50		500	X	100	1000	1		X	X				200	X	50	
6041 RS 152	A4868/76	MARBLE	TIDNAMURKUNA VOLCANICS	X	X	X	500	X	300	100	30		500	X	300	1000	10		X	X					X	50	
6041 RS 153	A4869/76	AMPHIBOLITE	BALTUCOODNA QUARTZITE	X	X	X	X	2	20	80	80		3000	10	100	500	5		X	X				300	X	100	
6041 RS 154	A4870/76	RHYOLITE	TIDNAMURKUNA VOLCANICS	X	X	X	300	X	80	70	200		100	X	80	1000	10		X	X					X	100	
6041 RS 155	A4871/76	"	"	X	X	X	700	X	5	80	100		100	10	100	X	20		X	2					X	X	
6041 RS 156	A4872/76	EPIDOSITE	"	X	X	X	1500	X	5	50	10		500	10	70	500	20		X	1					X	X	
6041 RS 157	A4858/76	DIORITE	DYKE;	X	X	X	300	X	200	300	200		800	X	300	500	3		X	X				500	X	70	
6041 RS 161	A4881/76	BASALT	WIRRIE CURRIE GRANITE			+	390		70	280	25	40		+	320		30	110				140	N.D.	N.D.	300	55	80
6041 RS 162	A4882/76	"	CADLAREENA VOLCANICS			+	90		50	520	20	45		10	160		12	36				150	N.D.	N.D.	320	45	75
6041 RS 163	A4883/76	"	"			+	260		55	400	10	55		+	120		14	46				100	N.D.	N.D.	330	40	110
6041 RS 164	A4884/76	"	"			0.05	70		50	280	30	20		10	190		12	28				260	N.D.	N.D.	280	35	85
6042 RS 1	A4846/76	EPIDOSITE	UNNAMED METAMORPHICS Ed <sub>1</sub>	X	X	X	500	X	150	100	100		1000	X	300	500	10		X	X				300	X	50	
6140 RS 4	A555/62	SYENITE	CADLAREENA VOLCANICS	0.2		X			30	1000	15					120		100						700		80	
6140 RS 5	A556/62	"	"	0.2		X			30	400	3					70		30						600		40	
6140 RS 6	A557/62	"	"	0.1		X			25	500	20					80		20						600		25	
6140 RS 7	A558/62	"	"	0.2		X			25	300	3					100		15						500		30	
6140 RS 8	A559/62	"	"	0.2		X			70	400	15					120		20						600		70	
6140 RS 15	A4838/76	QUARTZ-CROSSITE ROCK	DUFF CREEK BEDS	X	X	X	X	X	30	50	10 000		10 000	10	80	200	3		X	2				80	X	300	
6140 RS 16	A4839/76	BIOTITE-CROSSITE ROCK	"	X	X	X	X	X	10	80	300		2000	X	70	1000	1		X	1				80	X	200	
6140 RS 17	A4840/76	SPHERULITES	BUNGADILLINA MONZONITE	X	X	X	1500	X	30	100	300		300	X	80	10 000	10		X	1				200	X	X	