

SOUTH AUSTRALIA  
DEPARTMENT OF MINES AND ENERGY



**OPEN FILE ENVELOPE NO. 5183**

**PEL 16**

**MURRAY BASIN**

**OVERLAND CORNER 1**

**TEST REPORTS**

Submitted by

**SADME**

1986

SCANNED

(c) South Australian Department of Mines and Energy: 17/12/92

This report was supplied as part of the requirement to hold a mineral or petroleum exploration tenement in the State of South Australia. The Department accepts no responsibility for statements made, or conclusions drawn, in the report or for the quality of original text or drawings.

All rights reserved under the copyright. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the written permission of the S.A. Department of Mines and Energy, P.O. Box 151, Eastwood, S.A. 5063.

**ENVELOPE 5183**

**TENEMENT:** PEL 16; Murray Basin

**TENEMENT HOLDER:** International Mining Corporation NL (operator) and Satina Pty Ltd

**CONTENTS**

	<b>SADME NO.</b>
<b>REPORTS:</b> Preiss, W.V., 1984. Micropalaeontological examination of drillcore from the basement rocks in Overland Corner (Murray Basin). <i>South Australia. Department of Mines and Energy. Report Book, number 830.</i>	Not microfilmed.
Webb, A.W., 1986. Evaluation of suitability for K-Ar and Rb-Sr dating. (Amdel report no. G6753/86, 7 July 1986).	<b>5183 R 1</b> Pgs 3-4

**END OF CONTENTS**

DEPARTMENT OF MINES AND ENERGY  
SOUTH AUSTRALIA

OPEN FILE	
21/12/92	<i>dlg</i>
Date	Initials
Released	

REPT.BK.NO. 830  
MICROPALAEONTOLOGICAL EXAMINATION  
OF DRILLCORE FROM THE BASEMENT  
ROCKS IN OVERLAND CORNER 1  
(MURRAY BASIN)

by

W.V. PREISS

AUGUST, 1984

SR.27/4/478  
Biostrat.No.10/84



<u>CONTENTS</u>	<u>PAGE</u>
ABSTRACT	1
LOCATION AND GEOLOGY OF OVERLAND CORNER 1	1
DESCRIPTION OF DRILLCORE	2
EXAMINATION OF MACERATIONS	3
CONCLUSIONS AND RECOMMENDATIONS	3
REFERENCES	5

DEPARTMENT OF MINES AND ENERGY  
SOUTH AUSTRALIA

REPT. BK. NO. 830  
DME NO. SR 27/4/478  
BIOSTRAT. NO. 10/84  
DISK NO. 11

MICROPALAEONTOLOGICAL EXAMINATION OF DRILLCORE FROM  
THE BASEMENT ROCKS IN OVERLAND CORNER 1  
(MURRAY BASIN)

ABSTRACT

The Overland Corner 1 petroleum exploration well, Canegrass Lobe, Murray Basin, entered 12 m of steeply dipping grey siltstone as 'basement' below Permo-Carboniferous, Mesozoic and Tertiary strata. The siltstone contains fine sandy laminations and small carbonaceous shale flakes. There are spaced fractures but no penetrative slaty cleavage, and, surprisingly, no visible metamorphism. No indigenous microfossils were found in either thin section or maceration, and there is no biostratigraphic evidence of the age.

INTRODUCTION

Micropalaeontological examination of drillcore from the 'basement' rocks encountered in International Mining Corporation Overland Corner 1 (Murray Basin) follows from interest expressed by the company in the economic potential of these rocks, and in particular, whether they are unproductive Kanmantoo Group (normally metamorphosed) or mid-Palaeozoic strata that may have some hydrocarbon potential.

LOCATION AND GEOLOGY OF OVERLAND CORNER 1

The Overland Corner 1 drillsite is situated 10 km north of Overland Corner at Lat. 34°04'06"S, Long. 140°18'03"E, within the Canegrass Lobe in the western Murray Basin, immediately west of the Hamley Fault as shown on the RENMARK 1:250 000 sheet (Firman, 1972).

The drillhole penetrated Tertiary, Mesozoic and Permo-Carboniferous strata to a depth of 642.1 m, and then intersected some 12 m of 'basement' rocks to a total depth of 654.7 m. The nearest other drillholes that have entered rocks older than late Palaeozoic are Berri North 1 (34 km ESE of Overland Corner 1),

which entered metagreywacke, and Loxton 2 (60 km SSE of Overland Corner 1), which entered greywacke (Thornton, 1974). Other bores south of the River Murray entered metasediments that have been correlated with the Kanmantoo Group (Thornton, 1974), as have the greywackes in Berri North 1 and Loxton 2. However, on the CHOWILLA sheet to the north, Rogers (1977), suggested that the basement north of and perhaps beneath the Canegrass Lobe is largely Adelaidean, but drillhole data are very sparse. Thornton (1974) discussed the mainly arenaceous sediments, of possible Devonian age, in the Renmark Trough but expressed uncertainty as to the western extent of these sediments in the Canegrass Lobe. Thus available data suggest that the pre-Permo-Carboniferous 'basement' in Overland Corner 1 could be Adelaidean, Cambrian Kanmantoo Group, or (?) Devonian.

#### Description of Drillcore

The 12 m core consists of rather uniform, medium grey siltstone which appears to be practically unmetamorphosed. A bedding lamination, a few millimetres thick, is expressed by grain size variations, and small wispy, dark grey flakes of very fine grained material are scattered parallel to the bedding. The bedding dips at about 70° (assuming vertical drillhole) and there is an incipient fracture cleavage dipping at about 45° in the opposite direction. No penetrative slaty cleavage is visible.

In thin section (T.S. 45420, from 643.7 m) the rock consists of alternating, crudely defined bands of siltstone and fine sandstone. The siltstone consists of very fine grained quartz, clay minerals and fine flakes of white mica, the latter being almost completely randomly oriented. Locally, there is a slight tendency to parallelism with bedding. The sandy laminae are 1 to 2 mm thick and contain angular to subangular quartz and minor carbonate, up to 0.1 mm in grain size, scattered in a silty matrix. Their boundaries are rather diffuse. The silty and sandy laminations appear to contain little or no organic matter. However, the siltstone does include wispy fragments of carbonaceous shale which lack quartz and mica flakes; their pale greyish-brown colour in thin section suggests that these do contain organic matter. The rock is cut by a number of spaced fractures at a high angle to bedding, and limonite staining along

these fractures indicates that the rock has been subject to slight pre-Permo-Carboniferous weathering. No microfossils were found in thin section, either in the siltstone or in the shale clasts.

#### EXAMINATION OF MACERATIONS

Samples of grey siltstone were collected from 654.5 m and 643.7 m. These were macerated in hot HF and hot concentrated HCl, and temporary slides of the washed residues were examined. These contained pale greyish-brown (?) organic material presumed to have originated from the carbonaceous shale clasts; it was mostly clumped into irregular masses which were difficult to disperse, and in which no evidence of microfossils was seen. Further treatment by light oxidation with Schultz solution and heavy liquid separation of the remaining mineral material produced moderate quantities of residue of brownish-grey, structureless organic material. No indigenous microfossils were found, although, in two slides, single, poorly preserved specimens of the Cretaceous spores Cicatricosisporites australiensis and ?Todisporites sp. occur (W.K. Harris, pers. comm., July, 1984). Leitz Orthoplan stage coordinates are 464-941 (slide S5942-1) and 581-991 (slide S5941-1). Both samples are from 643.7 m, just below the late Palaeozoic unconformity. These spores are clearly contaminants, probably from drilling mud adhering to the core.

#### CONCLUSIONS AND RECOMMENDATIONS

The steeply dipping siltstone 'basement' in Overland Corner 1 contains no indigenous microfossils. There is therefore no biostratigraphic evidence for the age of these rocks. Lithologically, they resemble a number of late Proterozoic formations known from the Adelaide Geosyncline, rather than Kanmantoo Group. The apparent absence of even low grade metamorphism is surprising, however. Although there is no evidence of a mid-Palaeozoic age for these rocks, and their steep dip strongly suggests that they are pre-Delamerian, they do contain organic material of moderately light colour. It is

recommended that the thermal maturation and total organic carbon be determined before they can be ruled out as potential source rocks for the generation of petroleum or gas.

*W.V. Preiss.*

W.V. Preiss

Biostratigraphy Branch



## REFERENCES

- Firman, J.B. (Compiler), 1972. RENMARK, South Australia. Explanatory Notes, 1:250 000 geological series. Sheet SI/54-10. Geol. Surv. S. Aust.
- Rogers, P.A. (Compiler), 1977. CHOWILLA, South Australia. Explanatory Notes, 1:250 000 geological series. Sheet SI/54-6. Geol. Surv. S. Aust.
- Thornton, R.C.N., 1974. Hydrocarbon potential of western Murray Basin and infrabasins. Rep. Invest., geol. Surv. S. Aust., 41.



**The Australian  
Mineral Development  
Laboratories**

Flemington Street, Frewville,  
South Australia 5063  
Phone Adelaide (08) 79 1662  
Telex AA82520

Please address all  
correspondence to  
P.O. Box 114 Eastwood  
SA 5063  
In reply quote:

# amdel

00003

7 July 1986

GS 1/2/0

OPEN FILE	
21/12/92	dlg
Date	Initials
Released	

South Australian Department of  
Mines & Energy,  
P.O. Box 151,  
EASTWOOD, S.A. 5063

ATT: DR. D. GRAVESTOCK

REPORT G 6753/86

YOUR REFERENCE:	11/06/0800. EX-541
IDENTIFICATION:	A907/86
MATERIAL:	Core
LOCALITY:	Overland Corner No. 1, 652.6-652.8m
DATE RECEIVED:	17 June 1986
WORK REQUIRED:	Evaluation of suitability for K-Ar and Rb-Sr dating.

Investigation and Report by: Dr Alan W Webb

Manager - Geological Services Section: Dr Keith J Henley

Head Office:  
Flemington Street, Frewville  
South Australia 5063  
Telephone (08) 79 1662  
Telex: Amdel AA82520  
Pilot Plant:  
Osman Place  
Thebarton, S.A.  
Telephone (08) 43 5733  
Telex: Amdel AA82725  
Branch Laboratories:  
Melbourne, Vic.  
Telephone (03) 645 3093  
Perth, W.A.  
Telephone (09) 325 7311  
Telex: Amdel AA94893  
Sydney, N.S.W.  
Telephone (02) 439 7735  
Telex: Amdel AA20053  
Townsville  
Queensland 4814  
Telephone (077) 75 1377

*Keith Henley*

for Dr William G Spencer  
General Manager  
Applied Sciences Group

bp

EVALUATION OF SUITABILITY FOR DATING OF ONE SAMPLE OF DRILL CORE

## 1. INTRODUCTION

One sample of drill core (A907/86) from Overland Corner No. 1 was received from Dr. D. Gravestock, South Australian Department of Mines & Energy, with a request to carry out an evaluation of suitability for isotopic dating by either the K-Ar or Rb-Sr methods.

## 2. EVALUATION

A thin section (TSC47512) was prepared and examined. The rock is a calcareous siltstone or silty shale with detrital quartz and micas (mainly very fine flakes of muscovite, with minor biotite).

The detrital mica component may not be a serious impediment to dating since the most appropriate method would be to separate the -2 micron size fraction for dating, thereby eliminating most of the detrital flakes. However, it may prove very difficult to remove the significant carbonate phase which would interfere with both K-Ar and Rb-Sr dating.

If Rb-Sr dating were to be attempted, it would be necessary to analyse a minimum of 4 samples from the same unit.

At this stage, because the petrographic evidence casts doubt on the suitability for Rb-Sr dating, the approximate Rb and Sr concentrations normally determined under Code M2.1 have not been made.